For Research Use Only

FBXO22 Polyclonal antibody

Catalog Number: 13606-1-AP

Featured Product

20 Publications

BC039024



Purification Method:

WB 1:500-1:4000

IHC 1:50-1:500

for WB

Antigen affinity purification

IP 0.5-4.0 ug for IP and 1:500-1:1000

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number:

GeneID (NCBI):

150ul, Concentration: 500 µg/ml by 26263

Nanodrop; Full Name:

Source: F-box protein 22 Rabbit Calculated MW: Isotype: 403 aa, 45 kDa

IgG Observed MW: Immunogen Catalog Number: 45 kDa

AG4485

13606-1-AP

Positive Controls:

WB: HeLa cells, HuH-7 cells, HepG2 cells

IP: HeLa cells,

IHC: mouse heart tissue,

Applications

Tested Applications: IHC, IP, WB, ELISA

Cited Applications: IHC, IP, WB

Species Specificity: human, mouse

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

FBXO22, also named as FBX22, is a substrate-recognition of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex. FBXO22 is a key regulator of senescence induction through ubiquitylation of p53 for degradation. FBXO22 also acts as a molecular switch for the antagonistic and agonistic actions of selective estrogen receptor modulators (SERM) and determines the sensitivity of breast cancer to SERM by ubiquitylating KDM4B complexed with unliganded or SERMs-bound estrogen receptor (ER). (PMID: 32536008, PMID: 32249768)

Notable Publications

Author	Pubmed ID	Journal	Application
Min Lin	36127346	Cell Death Dis	WB,IHC
Siqiaozhi Li	36112263	Cell Biol Toxicol	IHC
Rita Moreno	36084789	Free Radic Biol Med	WB

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 13606-1-AP (FBXO22 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



IP Result of anti-FBXO 22 (IP:13606-1-AP, 4ug; Detection:13606-1-AP 1:500) with HeLa cells lysate 4000ug.



Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 13606-1-AP (FBXO22 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).