For Research Use Only

IL3RA/CD123 Polyclonal antibody

Catalog Number: 13655-1-AP



Basic Information

Catalog Number:

13655-1-AP

GenBank Accession Number:

BC035407

UNIPROT ID:

Full Name:

P26951

GeneID (NCBI): Size:

150ul, Concentration: 200 ug/ml by

Nanodrop and 133 ug/ml by Bradford ENSEMBL Gene ID:

method using BSA as the standard; ENSG00000185291 Source:

Rabbit Isotype:

interleukin 3 receptor, alpha (low

Immunogen Catalog Number: affinity)

AG4452 Calculated MW:

> 378 aa. 43 kDa Observed MW: 70 kDa

Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:2400 IHC 1:50-1:500

Applications

Tested Applications:

WB, IHC, ELISA

Species Specificity:

Positive Controls:

WB: HL-60 cells,

IHC: human tonsillitis tissue,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Interleukin-3 receptor subunit alpha (IL3RA), also known as CD123, belongs to the cytokine receptor family. It is a glycoprotein containing an extracellular domain, involving a predicted Ig-like domain, two FnIII domains, a transmembrane domain and an intracellular domain (PMID: 24513123). IL3RA is expressed on hematopoietic progenitor cells, plasmacytoid dendritic cells, basophils, monocytes, some B cells, and in a variety of hematologic malignancies (PMID: 10527461; 34855461). IL-3 initially binds to IL3RA and subsequently recruits the beta chain (CD131) to form the high-affinity receptor, resulting in the activation of a series of signaling pathways including the JAK/STAT, Ras-MAPK, and phosphatidylinositol 3-kinase pathways (PMID: 29296749). IL-3R signaling regulates the proliferation, survival, and differentiation of hematopoietic cells.

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

*** 20ul sizes contain 0.1% BSA

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



HL-60 cells were subjected to SDS PAGE followed by western blot with 13655-1-AP (IL3RA antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 13655-1-AP (IL3RA antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 13655-1-AP (IL3RA antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).