## For Research Use Only

# FGF12 Polyclonal antibody

Catalog Number: 13784-1-AP

Featured Product

5 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number:

13784-1-AP BC022524
Size: Genel D (NCBI):

150ul , Concentration: 260 ug/ml by 2257

Nanodrop; UNIPROT ID:
Source: P61328
Rabbit Full Name:

Isotype: fibroblast growth factor 12

IgG Calculated MW:
Immunogen Catalog Number: 181 aa, 20 kDa
AG4797 Observed MW:
20-22 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:2000-1:12000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:20-1:200

**Applications** 

Tested Applications: WB, IP, IHC, ELISA Cited Applications:

WB, IHC, IF Species Specificity: human, mouse, rat

Cited Species: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### **Positive Controls:**

WB: mouse brain tissue, rat brain tissue

IP: mouse brain tissue,
IHC: mouse brain tissue,

# **Background Information**

FGF 12 is a member of the FGF family. FGF family members play important roles in embryogenesis, angiogenesis, and wound repair. FGF 12 lacks the N-terminal signal sequence present in most of the FGF family members, but it contains clusters of basic residues that have been demonstrated to act as a nuclear localization signal. When transfected into mammalian cells, this protein accumulated in the nucleus, but was not secreted. FGF 12 plays an intracellular role in the inhibition of radiation-induced apoptosis. FGF 12 is expressed abundantly in developing and adult nervous systems; therefore, FGF 12 was thought to be related to nervous system development and function.

### **Notable Publications**

Author	Pubmed ID	Journal	Application
Jana Velíšková	33982289	Epilepsia	WB
Yukiko Hanada	30065296	Sci Rep	WB,IF
Yudong Gao	39122707	Nat Commun	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) APTGLAB (1-888-478-4522) (foll free F: proteintech@ptglab.com


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

# **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 13784-1-AP (FGF 12 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



IP result of anti-FGF12 (IP:13784-1-AP, 4ug; Detection:13784-1-AP 1:6000) with mouse brain tissue lysate 1280 ug.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 13784-1-AP (FGF12 Antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 13784-1-AP (FGF12 Antibody) at dilution of 1:50 (under 40x lens).