

For Research Use Only

TACC1 Polyclonal antibody

Catalog Number: 13862-1-AP **1 Publications**



Basic Information

Catalog Number: 13862-1-AP	GenBank Accession Number: BC041391	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 900 µg/ml by Nanodrop and 387 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 6867	Recommended Dilutions: WB 1:500-1:2000 IHC 1:250-1:1000 IF 1:20-1:200
Source: Rabbit	Full Name: transforming, acidic coiled-coil containing protein 1	
Isotype: IgG	Calculated MW: 805 aa, 88 kDa	
Immunogen Catalog Number: AG4842	Observed MW: 110-130 kDa	

Applications

Tested Applications: IF, IHC, WB, ELISA	Positive Controls: WB : MCF7 cells, HeLa cells IHC : human stomach cancer tissue, IF : HeLa cells,
Cited Applications: IF	
Species Specificity: human	
Cited Species: human	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Pan Xu	34897285	Cell Death Discov	IF

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




MCF7 cells were subjected to SDS PAGE followed by western blot with 13862-1-AP (TACC1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue slide using 13862-1-AP (TACC1 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue slide using 13862-1-AP (TACC1 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of HeLa cells using 13862-1-AP (TACC1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).