

For Research Use Only

ZEB2 Polyclonal antibody

Catalog Number: 14026-1-AP

Featured Product

58 Publications



Basic Information

Catalog Number: 14026-1-AP	GenBank Accession Number: BC060819	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 850 µg/ml by Nanodrop;	GeneID (NCBI): 9839	Recommended Dilutions: WB 1:2000-1:12000 IHC 1:50-1:500
Source: Rabbit	UNIPROT ID: O60315	
Isotype: IgG	Full Name: zinc finger E-box binding homeobox 2	
Immunogen Catalog Number: AG5171	Calculated MW: 136 kDa	
	Observed MW: 150-170 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls: WB : A549 cells, HepG2 cells, HeLa cells
Cited Applications: ChIP, IF, IHC, WB	IHC : human gliomas tissue, human lung cancer tissue, human renal cell carcinoma tissue
Species Specificity: human	
Cited Species: human, chicken, rat, mouse, zebrafish	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

Zinc finger E-box-binding homeobox (ZEB2) is a key transcription factor that acts as a multifunctional regulator during nervous system development. ZEB2 contains two zinc finger domains and a homeodomain-like sequence and interacts with the TGF- β superfamily signaling regulators, Smads, to regulate the expression of their downstream genes. ZEB2 is expressed in the developing neural tube, as well as in neural crest cells, the hippocampus and the cerebral cortex. ZEB2 has previously been implicated in EMT, cell-cycle progression, apoptosis and senescence. ZEB2 was overexpressed in bladder, ovarian, stomach, pancreatic and squamous cell carcinoma, in the intestinal subtype of stomach cancers, and at the invasive front of CRC where EMT is most prominent. ZEB2 also mediates cell-fate decision in neuronal, T cells and hematopoietic stem cells. The calculated molecular weight of ZEB2 is 136 kDa, but we find the 95 kDa band in some publication (PMID: 27659015)

Notable Publications

Author	Pubmed ID	Journal	Application
Liming Wang	31566718	J Cell Physiol	WB
Rongkun Li	34568024	Front Oncol	WB
Mengying Li	34492118	J Periodontal Res	WB,IHC

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 14026-1-AP (ZEB2 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 14026-1-AP (ZEB2 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




WB result of ZEB2 antibody (14026-1-AP; 1:6000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ZEB2 transfected HepG2 cells.



Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 14026-1-AP (ZEB2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human renal cell carcinoma tissue slide using 14026-1-AP (ZEB2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human gliomas tissue slide using 14026-1-AP (ZEB2 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).