For Research Use Only

NINJ2 Polyclonal antibody

Catalog Number: 14085-1-AP



Basic Information

Catalog Number: 14085-1-AP

GenBank Accession Number:

Purification Method: Antigen affinity purification

Size:

GeneID (NCBI):

Recommended Dilutions:

IHC 1:50-1:500

150ul , Concentration: 300 $\mu g/ml$ by Nanodrop and 213 µg/ml by Bradford Full Name:

4815

BC057766

niniurin 2

method using BSA as the standard;

Calculated MW:

Rabbit

142 aa, 16 kDa

Isotype: IgG

Immunogen Catalog Number:

AG5215

Applications

Tested Applications:

IHC,ELISA

Positive Controls:

Species Specificity:

IHC: human tonsillitis tissue,

human, mouse, rat Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Background Information

NINJ2 (Ninjurin2) is a member of the ninjurin family of adhesion molecules which mediate cell-to-cell and cell-toextracellular matrix interactions during development, differentiation, and regeneration of the peripheral nervous system. The gene encodes NINJ2 is located on chromosome 12p13. NINJ2 mRNA is widely expressed in adult human tissues, with highest level in bone marrow, followed by peripheral leukocytes, lung, and lymph nodes. In the peripheral nervous system, NINJ2 is expressed constitutively in mature sensory and enteric neurons. The expression of NINJ2 is upregulated after nerve injury in Schwann cells, suggesting that it may promote nerve regeneration. It may also play an important role in the pathogenesis of inflammatory disorder. (PMID: 10627596)

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 14085-1-AP (NINJ2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 14085-1-AP (NINJ2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).