

For Research Use Only

STAU1 Polyclonal antibody

Catalog Number: 14225-1-AP

Featured Product

21 Publications



Basic Information

Catalog Number:

14225-1-AP

Size:

150ul, Concentration: 500 ug/ml by Nanodrop and 333 ug/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG5464

GenBank Accession Number:

BC050432

GeneID (NCBI):

6780

UNIPROT ID:

O95793

Full Name:

staufen, RNA binding protein, homolog 1 (Drosophila)

Calculated MW:

63 kDa

Observed MW:

55 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:500-1:2000

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IHC, IF, RIP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, pig, monkey

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : human brain tissue, K-562 cells, mouse brain tissue

IP : mouse brain tissue,

IHC : mouse brain tissue, mouse pancreas tissue

IF/ICC : A549 cells,

Background Information

Staufen1 (STAU1)-mediated mRNA decay (SMD) degrades translationally active mRNAs that bind the double-stranded (ds)RNA binding protein STAU1 within their 3'-untranslated regions (3'UTRs) [PMID:21307942]. STAU1 can binds to stem structures and protect Dvl2 mRNA from degradation by associating with the secondary structure in the U1 region which is located at 5' position of second U stretch in the U2 region [PMID:17510634]. It has a role in specific positioning of mRNAs at given sites in the cell by cross-linking cytoskeletal and RNA components, and in stimulating their translation at the site. There are two isoforms of STAU1 protein and the molecular weight of two isoforms are respectively 63 kDa and 55 kDa

Notable Publications

Author	Pubmed ID	Journal	Application
Evan P Booy	30247708	Nucleic Acids Res	WB
Shuyuan Shen	32322670	Mol Ther Oncolytics	RIP
Yoichiro Sugimoto	25799984	Nature	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human brain tissue were subjected to SDS PAGE followed by western blot with 14225-1-AP (STAU1 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.




IP result of anti-STAU1 (IP:14225-1-AP, 4ug; Detection:14225-1-AP 1:500) with mouse brain tissue lysate 3440ug.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 14225-1-AP (STAU1 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse pancreas tissue slide using 14225-1-AP (STAU1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed A549 cells using STAU1 antibody (14225-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).