For Research Use Only

NCAM1/CD56 Polyclonal antibody

Catalog Number: 14255-1-AP

Featured Product

68 Publications



Basic Information

Catalog Number:

14255-1-AP

BC047244 GeneID (NCBI):

150ul , Concentration: 850 ug/ml by

Nanodrop: ENSEMBL Gene ID: ENSG00000149294 Rabbit **UNIPROT ID:**

Isotype P13591 IgG Full Name:

Immunogen Catalog Number: neural cell adhesion molecule 1

AG5528 Calculated MW:

95 kDa Observed MW:

120 kDa, 140 kDa, 180 kDa

GenBank Accession Number:

Purification Method: Antigen affinity purification Recommended Dilutions:

WB: 1:5000-1:50000 IHC: 1:8000-1:32000 IF/ICC: 1:50-1:500

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Cited Applications: WB, IHC, IF

Species Specificity: human, mouse, rat, pig

Cited Species:

human, mouse, rat, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse brain tissue, Neuro-2a cells, rat brain tissue, pig brain tissue

IHC: human tonsillitis tissue, human appendicitis tissue, human colon tissue, human lung cancer tissue, Insulinoma tissue, mouse brain tissue, mouse colon tissue, rat brain tissue

IF/ICC: SH-SY5Y cells.

Background Information

Neural cell adhesion molecule 1 (NCAM1, also known as CD56) is a cell adhesion glycoprotein of the immunoglobulin (Ig) superfamily. It is a multifunction protein involved in synaptic plasticity, neurodevelopment, and neurogenesis. NCAM1 is expressed on human neurons, glial cells, skeletal muscle cells, NK cells and a subset of T cells, and the expression is observed in a wide variety of human tumors, including myeloma, myeloid leukemia, neuroendocrine tumors, Wilms' tumor, neuroblastoma, and NK/T cell lymphomas. Three major isoforms of NCAM1, with molecular masses of 120, 140, and 180 kDa, are generated by alternative splicing of mRNA (PMID: 9696812). The glycosylphosphatidylinositol (GPI)-anchored NCAM120 and the transmembrane NCAM140 and NCAM180 consist of five Ig-like domains and two fibronection-type III repeats (FNIII). All three forms can be posttranslationally modified by addition of polysialic acid (PSA) (PMID: 14976519). Several other isofroms have also been described (PMID: 1856291).

Notable Publications

Author	Pubmed ID	Journal	Application
Yanping Li	36213822	J Oncol	IF
Shaolong Li	30264546	Cancer Sci	IHC
Ashley Gillon	26385499	Biogerontology	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20° C storage


*** 20ul sizes contain 0.1% BSA

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




WB result of NCAM1/CD56 antibody (14255-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-NCAM1/CD56 transfected Neuro-2a cells.




Various lysates were subjected to SDS PAGE followed by western blot with 14255-1-AP (NCAM1/CD56 antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 14255-1-AP (NCAM1/CD56 antibody) at dilution of 1:16000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 14255-1-AP (NCAM1/CD56 antibody) at dilution of 1:16000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed SH-SY5Y cells using NCAM1/CD56 antibody (14255-1-AP) at dilution of 1:200 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).