

For Research Use Only

Fascin Polyclonal antibody

Catalog Number: 14384-1-AP

Featured Product

11 Publications



Basic Information

Catalog Number: 14384-1-AP	GenBank Accession Number: BC000521	Purification Method: Antigen affinity purification
Size: 150ul, Concentration: 400 ug/ml by Nanodrop and 300 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 6624	Recommended Dilutions: WB 1:2000-1:8000 IHC 1:50-1:500 IF-P 1:50-1:500
Source: Rabbit	ENSEMBL Gene ID: ENSG00000075618	
Isotype: IgG	UNIPROT ID: Q16658	
Immunogen Catalog Number: AG5816	Full Name: fascin homolog 1, actin-bundling protein (Strongylocentrotus purpuratus)	
	Calculated MW: 55 kDa	
	Observed MW: 50-55 kDa	

Applications

Tested Applications: WB, IHC, IF-P, FC (Intra), ELISA	Positive Controls: WB: HeLa cells, mouse thymus tissue, K-562 cells, NIH/3T3 cells, SH-SY5Y cells IHC: human tonsillitis tissue, human liver cancer tissue IF-P: mouse brain tissue,
Cited Applications: WB, IHC, IF	
Species Specificity: human, mouse	
Cited Species: human, mouse, rat	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Fascin is a highly conserved and ubiquitous actin cross-linking protein that has a major function in cell motility and adhesion. Fascin localizes to a number of highly dynamic cellular structures that require strong mechanical support which include stress fibers and cellular protrusions such as microvilli, microspikes and lamellipodia. Fascin is up-regulated in many human carcinomas. It's a marker for Reed-Sternberg cells in Hodgkin's disease; it is also a highly selective marker for dendritic cells of lymphoid tissues.

Notable Publications

Author	Pubmed ID	Journal	Application
Lihua Guo	25433493	J Exp Clin Cancer Res	WB
Peiyu Gao	32554050	Immunol Lett	WB
Ajab Khan	31465800	Int J Biol Macromol	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 14384-1-AP (Fascin antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 14384-1-AP (Fascin antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 14384-1-AP (Fascin antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



WB result of Fascin antibody (14384-1-AP; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-Fascin transfected HeLa cells.



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using 14384-1-AP (Fascin antibody) at dilution of 1:50 and Alexa Fluor 488-Conjugated Goat Anti-Rabbit IgG(H+L).



1×10^6 SH-SY5Y cells were intracellularly stained with 0.2 ug Anti-Human Fascin (14384-1-AP) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with 0.1% TritonX-100.