

CYP17A1 Polyclonal antibody

Catalog Number: 14447-1-AP

39 Publications

Basic Information

Catalog Number: 14447-1-AP	GenBank Accession Number: BC062997	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 700 µg/ml by Nanodrop;	GeneID (NCBI): 1586	Recommended Dilutions: WB 1:2000-1:12000 IHC 1:100-1:400 IF 1:20-1:200
Source: Rabbit	Full Name: cytochrome P450, family 17, subfamily A, polypeptide 1	
Isotype: IgG	Calculated MW: 57 kDa	
Immunogen Catalog Number: AG5899	Observed MW: 48-51 kDa	

Applications

Tested Applications: IF, IHC, WB, ELISA	Positive Controls:
Cited Applications: IF, IHC, WB	WB: A549 cells, Jurkat cells, rat brain tissue, HeLa cells, HepG2 cells, mouse adrenal gland tissue, mouse brain tissue
Species Specificity: human, mouse, rat	IHC: human prostate cancer tissue,
Cited Species: human, rat, mouse	IF: HepG2 cells,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

The human cytochrome P450c17 enzyme is encoded by the CYP17A1 gene. It is the qualitative regulator of steroidogenesis in the human adrenals. In its absence, aldosterone is produced in the human adrenal zona glomerulosa. The CYP17A1 gene is expressed in the adrenal cortex and the gonads but not in the placenta (PMID:19403566). Although CYP17A1 has been shown to be a 50-kDa protein, the presence of a lower molecular mass (30 kDa) immunoreactive form in rat Leydig cells was previously shown and it is either a degradation product or a truncated form of the 50 kDa enzyme (PMID:15761033). Defects in CYP17A1 are the cause of adrenal hyperplasia type 5 (AH5) and CYP17A1 is an important target for inhibition in the treatment of prostate cancer (PMID 27505042, PMID 27505042).

Notable Publications

Author	Pubmed ID	Journal	Application
Yanyan Zhao	36152347	Biochem Biophys Res Commun	IHC
Yan Peng	33109277	Biol Res	WB
Jun Li	36338128	Front Bioeng Biotechnol	WB, IF

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
 Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
 T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
 E: proteintech@ptglab.com
 W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 14447-1-AP (CYP17A1 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of HepG2 cells, using CYP17A1 antibody 14447-1-AP at 1:50 dilution and Rhodamine-labeled goat anti-rabbit IgG (red). Blue pseudocolor = DAPI (fluorescent DNA dye).



Immunohistochemical analysis of paraffin-embedded human prostate cancer tissue slide using 14447-1-AP (CYP17A1 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human prostate cancer tissue slide using 14447-1-AP (CYP17A1 Antibody) at dilution of 1:200 (under 40x lens).