For Research Use Only

RABGGTA Polyclonal antibody

Catalog Number:14448-1-AP

Featured Product

3 Publications


Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: 14448-1-AP	GenBank Accession Num BC003093	iber:	Purification Method: Antigen affinity purification
	Size: 150ul , Concentration: 450 ug/ml by Nanodrop and 453 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI):		Recommended Dilutions:
		5875		WB 1:1000-1:4000 IHC 1:20-1:200
		JNIPROT ID: Incl. 120-11200 Q92696 IF/ICC 1:50-1:500		
	Source:	Full Name:		
	Rabbit Isotype: IgG Immunogen Catalog Number: AG5923	Rab geranylgeranyltransferase, alpha subunit Calculated MW:		
		Observed MW:		
		60 kDa		
		Applications	Tested Applications:	Positive Controls:
WB, IHC, IF/ICC, ELISA	WB : HEK-293 cells, Jurkat cells			
Cited Applications: WB, IF	IHC : human lung tissue, human brain tissue			
Species Specificity: human, mouse	IF/ICC : HepG2 cells,			
Cited Species: human, mouse				
	Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0	vely, antigen		
Background Information	TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0 GGTase II (RAB geranylgeranyl transf	rely, antigen ith citrate erase, RABGGTase) is a he perfamily of small monor membrane-bound organe	meric GTPase elles and vesi	s. RABGGTase is extensively involved icles and specifically catalyses the
	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0 GGTase II (RAB geranylgeranyl transf subunits, which belongs to the Ras su intracellular membrane trafficking of covalent attachment of GGPP to the co 11136552, 10737774, 35961942).	rely, antigen ith citrate erase, RABGGTase) is a he perfamily of small monor membrane-bound organe	meric GTPase elles and vesi	s. RABGGTase is extensively involved icles and specifically catalyses the
	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0GGTase II (RAB geranylgeranyl transf subunits, which belongs to the Ras su intracellular membrane trafficking of covalent attachment of GGPP to the co 11136552, 10737774, 35961942).Author	rely, antigen ith citrate erase, RABGGTase) is a he perfamily of small monor membrane-bound organo ysteine of a carboxyl-tern	meric GTPase elles and vesi ninal CC or C)	s. RABGGTase is extensively involved icles and specifically catalyses the KC motif of RAB proteins (PMID
Background Information	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0GGTase II (RAB geranylgeranyl transf subunits, which belongs to the Ras su intracellular membrane trafficking of covalent attachment of GGPP to the col 11136552, 10737774, 35961942).AuthorPub Liang Wei Wang315	rely, antigen ith citrate erase, RABGGTase) is a he perfamily of small monor imembrane-bound organ ysteine of a carboxyl-term med ID Journal	meric GTPase elles and vesi ninal CC or C	s. RABGGTase is extensively involved icles and specifically catalyses the KC motif of RAB proteins (PMID Application
	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0GGTase II (RAB geranylgeranyl transf subunits, which belongs to the Ras su intracellular membrane trafficking of covalent attachment of GGPP to the colspan="2">to the Ras su intracellular membrane trafficking of covalent attachment of GGPP to the colspan="2">to the Colspan="2">to the Ras su intracellular membrane trafficking of covalent attachment of GGPP to the colspan="2">to the Colspan="2" to the Colspan="	rely, antigen ith citrate erase, RABGGTase) is a he perfamily of small monor membrane-bound organ ysteine of a carboxyl-term med ID Journal 18366 PLoS Pat	meric GTPase elles and vesi ninal CC or C thog ath Dis	s. RABGGTase is extensively involved icles and specifically catalyses the KC motif of RAB proteins (PMID Application WB
	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0GGTase II (RAB geranylgeranyl transf subunits, which belongs to the Ras su intracellular membrane trafficking of covalent attachment of GGPP to the colspan="2">to the Ras su intracellular membrane trafficking of covalent attachment of GGPP to the colspan="2">to the Colspan="2">to the Ras su intracellular membrane trafficking of covalent attachment of GGPP to the colspan="2">to the Colspan="2" to the Colspan="	rely, antigen ith citrate erase, RABGGTase) is a he perfamily of small monor membrane-bound organi ysteine of a carboxyl-term med ID Journal 18366 PLoS Pat 46297 Cell Dea 42502 Brain Re er shipment.	meric GTPase elles and vesi ninal CC or C thog ath Dis	s. RABGGTase is extensively involved icles and specifically catalyses the KC motif of RAB proteins (PMID Application WB WB


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HEK-293 cells were subjected to SDS PAGE followed by western blot with 14448-1-AP (RABGGTA antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human lung using 14448-1-AP (RABGGTA antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human lung using 14448-1-AP (RABGGTA antibody) at dilution of 1:100 (under 40x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 14448-1-AP (RABGGTA antibody) at dilution of 1:50 and Alexa Fluor 488conjugated Goat Anti-Rabbit IgG(H+L).