For Research Use Only

TSG101 Polyclonal antibody

Catalog Number:14497-1-AP

Featured Product 229 Publications




Basic Information	Catalog Number: 14497-1-AP	GenBank Accession Number: BC002487	Purification Method: Antigen affinity purification			
	Size:	GenelD (NCBI):	Recommended Dilutions:			
	150ul, Concentration: 407 µg/ml by	7251	WB 1:1000-1:5000			
	Bradford method using BSA as the standard; Source: Rabbit	UNIPROT ID: Q99816 Full Name: tumor susceptibility gene 101	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:100-1:400 IF/ICC 1:10-1:100			
				Isotype: IgG	Calculated MW: 44 kDa	
				Immunogen Catalog Number: AG5920	Observed MW: 46 kDa	
	Applications	Tested Applications:	Positive Controls:			
		WB, IP, IF, IHC, ELISA	WB : NI	WB : NIH/3T3 cells, HeLa cells		
Cited Applications: WB, IHC, IF, IP, RIP		IP : HeL	a cells,			
Species Specificity:		IHC : human colon cancer tissue, human heart tiss				
human, mouse, rat		IF/ICC	: Hela cells,			
Cited Species: human, mouse, rat, pig, rabbit, canine, monkey, bovine						
Note-IHC: suggested antigen i TE buffer pH 9.0; (*) Alternati						
	retrieval may be performed w buffer pH 6.0					
Background Information	retrieval may be performed w buffer pH 6.0 TSG101(Tumor susceptibility gene 1 and the final stages of cytokinesis. It identified as a candidate tumor supp	v ith citrate 01 protein) is essential for endos plays a crucial role for cell proli ressor gene and belongs to the u	feration and cell survival. TSG101 has been biquitin-conjugating enzyme family. TSG101			
	retrieval may be performed w buffer pH 6.0 TSG101(Tumor susceptibility gene 1 and the final stages of cytokinesis. It identified as a candidate tumor supp a marker for exosome. This protein h and 32 kDa.	v ith citrate 01 protein) is essential for endos plays a crucial role for cell proli ressor gene and belongs to the u	feration and cell survival. TSG101 has been			
	retrieval may be performed w buffer pH 6.0 TSG101(Tumor susceptibility gene 1 and the final stages of cytokinesis. It identified as a candidate tumor supp a marker for exosome. This protein h and 32 kDa. Author Put	vith citrate 01 protein) is essential for endos plays a crucial role for cell proli ressor gene and belongs to the u as 2 isoforms produced by alterr	feration and cell survival. TSG 101 has been biquitin-conjugating enzyme family. TSG 101 hative splicing with the molecular mass of 44			
Background Information Notable Publications	retrieval may be performed w buffer pH 6.0 TSG101(Tumor susceptibility gene 1 and the final stages of cytokinesis. It identified as a candidate tumor supp a marker for exosome. This protein h and 32 kDa. Author Put Kristien Peeters 276	vith citrate 01 protein) is essential for endos plays a crucial role for cell proli ressor gene and belongs to the u as 2 isoforms produced by alterr omed ID Journal	biquitin-conjugating enzyme family. TSG101 hative splicing with the molecular mass of 44 Application			
	retrieval may be performed w buffer pH 6.0 TSG101(Tumor susceptibility gene 1 and the final stages of cytokinesis. It identified as a candidate tumor supp a marker for exosome. This protein h and 32 kDa. Author Put Kristien Peeters 276 Qiyun Yang 325	vith citrate 01 protein) is essential for endos plays a crucial role for cell proli ressor gene and belongs to the u as 2 isoforms produced by alterr omed ID Journal 686364 Ann Neurol	feration and cell survival. TSG 101 has been biquitin-conjugating enzyme family. TSG 101 hative splicing with the molecular mass of 44 Application WB			
	retrieval may be performed w buffer pH 6.0 TSG101(Tumor susceptibility gene 1 and the final stages of cytokinesis. It identified as a candidate tumor supp a marker for exosome. This protein h and 32 kDa. Author Put Kristien Peeters 276 Qiyun Yang 325	with citrate 01 protein) is essential for endos plays a crucial role for cell proli ressor gene and belongs to the u as 2 isoforms produced by alterr omed ID Journal 686364 Ann Neurol 986908 Andrologia 592398 Int J Pharm ter shipment.	feration and cell survival. TSC 101 has been biquitin-conjugating enzyme family. TSG 101 hative splicing with the molecular mass of 44 Application WB WB			


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 14497-1-AP (TSG101 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




IP result of anti-TSG101 (IP:14497-1-AP, 3ug; Detection:14497-1-AP 1:500) with HeLa cells lysate 1320ug.



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 14497-1-AP (TSG101 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 14497-1-AP (TSG101 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of Hela cells, using TSG101 antibody 14497-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).