

For Research Use Only

PSMB9 Polyclonal antibody

Catalog Number: 14544-1-AP

Featured Product

15 Publications



Basic Information

Catalog Number: 14544-1-AP	GenBank Accession Number: BC065513	Purification Method: Antigen affinity purification
Size: 150ul, Concentration: 400 µg/ml by Nanodrop and 233 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 5698	Recommended Dilutions: WB 1:1000-1:6000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:50-1:500 IF 1:50-1:500
Source: Rabbit	Full Name: proteasome (prosome, macropain) subunit, beta type, 9 (large multifunctional peptidase 2)	
Isotype: IgG	Calculated MW: 23 kDa	
Immunogen Catalog Number: AG6048	Observed MW: 20-23 kDa	

Applications

Tested Applications:
IF, IHC, IP, WB, ELISA

Cited Applications:
IF, IHC, WB

Species Specificity:
human, mouse, rat

Cited Species:
human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HL-60 cells, rat spleen tissue, mouse kidney tissue, mouse liver tissue, mouse lung tissue, mouse spleen tissue, rat liver tissue, Raji cells

IP: Raji cells,

IHC: human spleen tissue, human tonsillitis tissue, human renal cell carcinoma tissue, mouse bladder tissue, mouse stomach tissue, rat stomach tissue

IF: HepG2 cells,

Background Information

PSMB9, also named as LMP2, PSMB6i and RING12, belongs to the peptidase T1B family. The proteasome is a multicatalytic proteinase complex which is characterized by its ability to cleave peptides with Arg, Phe, Tyr, Leu, and Glu adjacent to the leaving group at neutral or slightly basic pH. Replacement of PSMB6 by PSMB9 increases the capacity of the immunoproteasome to cleave model peptides after hydrophobic and basic residues. High expression of PSMB9 associated with the poor prognosis of patients with NPC.

Notable Publications

Author	Pubmed ID	Journal	Application
Yuki Uchihara	35594857	Mol Cell	WB
Xin Xie	35710020	Biochem Pharmacol	WB
Xin Xie	32595507	Front Pharmacol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 14544-1-AP (PSMB9 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human spleen tissue slide using 14544-1-AP (PSMB9 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



mouse kidney tissue were subjected to SDS PAGE followed by western blot with 14544-1-AP (LMP2 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



IP Result of anti-LMP2 (IP:14544-1-AP, 4ug; Detection:14544-1-AP 1:300) with Raji cells lysate 2800ug.



Immunofluorescent analysis of (10% Formaldehyde) fixed HepG2 cells using 14544-1-AP (LMP2 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).