## For Research Use Only

## RINT1 Polyclonal antibody

Catalog Number: 14567-1-AP

1 Publications



**Basic Information** 

**Applications** 

Catalog Number: 14567-1-AP

GenBank Accession Number:

**Purification Method:** Antigen affinity purification

Size:

GeneID (NCBI):

Recommended Dilutions:

150ul , Concentration: 600  $\mu g/ml$  by Nanodrop and 327 µg/ml by Bradford Full Name:

BC068483

60561

WB 1:200-1:1000 IHC 1:20-1:200

RAD50 interactor 1

IF 1:10-1:100

method using BSA as the standard;

Calculated MW: 91 kDa

Rabbit Isotype:

Observed MW: 87 kDa

AG6145

IgG

Immunogen Catalog Number:

**Positive Controls:** 

IF THE WRITISA

**Cited Applications:** 

**Tested Applications:** 

IF

IHC: human breast cancer tissue, human liver cancer

Species Specificity: human, mouse, rat

**Cited Species:** 

human

IF: HepG2 cells,

WB: HepG2 cells,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

**Background Information** 

RINT-1 was first identified as a Rad50-interacting protein that participates in radiation-induced G2/M checkpoint  $control.\ It\ also\ involved\ in\ regulation\ of\ membrane\ traffic\ between\ the\ endoplasmic\ reticulum\ and\ the\ Golgi.\ It$ interacts with p130, an Rb family protein, and together they regulate the telomerase-independent telomere length. Otherwise, it interacts with Zeste white 10 with its N-terminal region throughout the cell cycle.

Notable Publications

Author Pubmed ID Journal Application Robert S Kiss 37774976 J Biol Chem

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## Selected Validation Data



HepG2 cells were subjected to SDS PAGE followed by western blot with 14567-1-AP (RINT1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human breast cancer using 14567-1-AP (RINT1 antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of HepG2 cells using 14567-1-AP (RINT1 antibody) at dilution of 1:25 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).