

For Research Use Only

HPS4 Polyclonal antibody

Catalog Number: 14627-1-AP

5 Publications



Basic Information

Catalog Number:

14627-1-AP

Size:

150ul, Concentration: 500 ug/ml by Nanodrop and 133 ug/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG6202

GenBank Accession Number:

BC065030

GeneID (NCBI):

89781

UNIPROT ID:

Q9NQG7

Full Name:

Hermansky-Pudlak syndrome 4

Calculated MW:

77 kDa

Observed MW:

70-90 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), IP, ELISA

Cited Applications:

WB

Species Specificity:

human

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: Jurkat cells, A375 cells, HeLa cells, K-562 cells

IP: HeLa cells, IP result of anti-HPS4 (14627-1-AP for IP and Detection) with HeLa cell lysate.

IHC: human liver tissue,

IF/ICC: HepG2 cells,

Background Information

Hermansky-Pudlak syndrome (HPS) is a genetic disease characterized by oculocutaneous albinism, bleeding due to platelet storage pool deficiency, and lysosomal storage defects. This syndrome results from defects of diverse cytoplasmic organelles including melanosomes, platelet dense granules and lysosomes. HPS1 and HPS4 are the most frequently mutated genes associated with HPS in humans. Both of HPS1 and HPS4 are components of two complexes involved in biogenesis of melanosome and lysosome-related organelles: BLOC-3 and BLOC-4. HPS4 is supposed to interact with HPS1 and stabilize HPS1. The human HPS4 migrates at about 90 kDa on SDS-PAGE, versus its predicated molecular mass of 77 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Seunghyi Kook	29190429	Am J Respir Cell Mol Biol	WB
Trieu-Duc Vu	35504437	Gene	WB
Joshi Stephen	28296950	PLoS One	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 14627-1-AP (HPS4 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 14627-1-AP (HPS4 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 14627-1-AP (HPS4 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).


IP & WB of 14627-1-AP with HeLa Cell



N/A.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using HPS4 antibody (14627-1-AP) at dilution of 1:400 and Coralite@488-Conjugated Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).



1×10^6 HepG2 cells were intracellularly stained with 0.4 ug HPS4 Polyclonal antibody (14627-1-AP) and Coralite@488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.4 ug Rabbit IgG control Rabbit PolyAb (30000-O-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).