For Research Use Only

DDX39B Polyclonal antibody

Catalog Number: 14798-1-AP

Featured Product

11 Publications



Basic Information

Catalog Number:

14798-1-AP

BC000361 GeneID (NCBI):

Positive Controls:

IP: Jurkat cells.

WB: Jurkat cells, rat spleen tissue

IHC: human breast cancer tissue,

Size:

150ul, Concentration: 213 µg/ml by Bradford method using BSA as the

7919

WB 1:500-1:2000

standard:

HLA-B associated transcript 1

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Rabbit Isotype:

IgG

Calculated MW: 53 kDa

Observed MW:

GenBank Accession Number:

Immunogen Catalog Number:

AG6512

48 kDa

Tested Applications:

human, mouse, rat

Cited Species:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen

buffer pH 6.0

Antigen affinity purification Recommended Dilutions:

Purification Method:

protein lysate IHC 1:50-1:500

Applications

IHC. IP. WB. FLISA

Cited Applications:

ChIP, IF, IHC, IP, WB

Species Specificity:

human, chicken, mouse, pig

retrieval may be performed with citrate

Background Information

DDX39B, is a 428 amino acid protein, which contains one helicase C-terminal domain, one helicase ATP-binding domain and belongs to the DEAD box helicase family. DECD subfamily. DDX39B localizes in the nucleus and cytoplasm. BAT1 is involved in nuclear export of spliced and unspliced mRNA. BAT1 as ATP dependent RNA helicase, may be a translation initiation factor and acts as a negative regulator of inflammatory cytokines. This antibody may cross-react with DDX39A.

Notable Publications

Author	Pubmed ID	Journal	Application
Qiu Tu	36109793	J Exp Clin Cancer Res	WB,IF
Lingcai Zhao	36084138	PLoS Pathog	IF,WB
Ashley Boehringer	29109432	Sci Rep	WB,IF,IP

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


*** 20ul sizes contain 0.1% BSA

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 14798-1-AP (UAP56 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Jurkat cells were subjected to SDS PAGE followed by western blot with 14798-1-AP (UAP56 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP Result of anti-UAP56 (IP:14798-1-AP, 4ug; Detection:14798-1-AP 1:500) with Jurkat cells lysate 1200ug.