

For Research Use Only

# EXOSC2 Polyclonal antibody

Catalog Number: 14805-1-AP

Featured Product

4 Publications



## Basic Information

<b>Catalog Number:</b> 14805-1-AP	<b>GenBank Accession Number:</b> BC000747	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul, Concentration: 350 µg/ml by Nanodrop and 213 µg/ml by Bradford method using BSA as the standard;	<b>GeneID (NCBI):</b> 23404	<b>Recommended Dilutions:</b> WB 1:500-1:2000 IP 0.5-4.0 ug for IP and 1:500-1:2000 for WB
<b>Source:</b> Rabbit	<b>Full Name:</b> exosome component 2	<b>IHC 1:20-1:200</b> <b>IF 1:20-1:200</b>
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 33 kDa	
<b>Immunogen Catalog Number:</b> AG6525	<b>Observed MW:</b> 30-33 kDa	

## Applications

### Tested Applications:

IF, IHC, IP, WB, ELISA

### Cited Applications:

IF, WB

### Species Specificity:

human

### Cited Species:

human

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

**WB:** HeLa cells, HepG2 cells, Jurkat cells, MCF-7 cells, HEK-293 cells

**IP:** HeLa cells,

**IHC:** human skin cancer tissue,

**IF:** MCF-7 cells,

## Background Information

In the nucleus, the RNA exosome complex is involved in proper maturation of stable RNA species such as rRNA, snoRNA and snRNA, in the elimination of RNA processing by-products and non-coding 'pervasive' transcripts, such as antisense RNA species and promoter-upstream transcripts (PROMPTS), and of mRNAs with processing defects, thereby limiting or excluding their export to the cytoplasm. In the cytoplasm, the RNA exosome complex is involved in general mRNA turnover and specifically degrades inherently unstable mRNAs containing AU-rich elements (AREs) within their 3' untranslated regions, and in RNA surveillance pathways, preventing translation of aberrant mRNAs [PMID:15346807]. EXOSC2 is a non-catalytic component of the RNA exosome complex that has 3'->5' exoribonuclease activity and involves in a multitude of cellular RNA processing and degradation events [PMID: 17545563].

## Notable Publications

Author	Pubmed ID	Journal	Application
Tobias Moll	36241425	Life Sci Alliance	WB
Hani Goodarzi	27259150	Cell	WB, IF
Jakob Trendel	30528433	Cell	

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




WB result of EXOSC2 antibody (14805-1-AP; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-EXOSC2 transfected HEK-293 cells.




HeLa cells were subjected to SDS PAGE followed by western blot with 14805-1-AP (EXOSC2 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human skin cancer using 14805-1-AP (EXOSC2 antibody) at dilution of 1:100 (under 40x lens).



Immunofluorescent analysis of MCF-7 cells, using EXOSC2 antibody 14805-1-AP at 1:50 dilution and Rhodamine-labeled goat anti-rabbit IgG (red). Blue pseudocolor = DAPI (fluorescent DNA dye).



IP Result of anti-EXOSC2 (IP:14805-1-AP, 4ug; Detection:14805-1-AP 1:1000) with HeLa cells lysate 1080ug.