

For Research Use Only

Arginase-2 Polyclonal antibody

Catalog Number: 14825-1-AP

Featured Product

10 Publications



Basic Information

Catalog Number:

14825-1-AP

Size:

150ul, Concentration: 700 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG6609

GenBank Accession Number:

BC001350

GeneID (NCBI):

384

UNIPROT ID:

P78540

Full Name:

arginase, type II

Calculated MW:

39 kDa

Observed MW:

39-42 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:4000

IHC 1:50-1:500

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

Positive Controls:

WB: Caco-2 cells, mouse kidney tissue, rat kidney tissue

IHC: human prostate cancer tissue, human kidney tissue

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Arginase 2 is composed of 354 amino acid residues, including an NH₂-terminal presequence for mitochondrial targeting and import. In the mitochondria, ornithine generated by Arginase 2 will give rise to glutamate via ornithine aminotransferase (OAT). Glutamate participates in several transamination reactions, including forming α -ketoglutarate (α KG) that may enter the TCA cycle and increase cycle intermediates and flux. Arginase 1 is mainly expressed in hepatocytes, and mice with a disruption of Arginase 1 gene die soon after birth. Arginase 2 is poorly expressed in hepatocytes, and most highly expressed in kidney, prostate, and immune cells such as monocyte/macrophages. (PMID: 25234945, PMID: 27214549)

Notable Publications

Author	Pubmed ID	Journal	Application
Hagai Tavori	25183802	J Lipid Res	WB
Lu Gao	30889486	Biomed Pharmacother	WB
Katherine M Halloran	35098299	Biol Reprod	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of ARG2 antibody (14825-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-Arginase-2 transfected Caco-2 cells.



Immunohistochemical analysis of paraffin-embedded human prostate cancer tissue slide using 14825-1-AP (ARG2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Caco-2 cells were subjected to SDS PAGE followed by western blot with 14825-1-AP (Arginase-2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 14825-1-AP (Arginase-2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.