

For Research Use Only

HSD17B7 Polyclonal antibody

Catalog Number: 14854-1-AP

Featured Product

5 Publications



Basic Information

Catalog Number:

14854-1-AP

Size:

150ul, Concentration: 500 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG6639

GenBank Accession Number:

BC065246

GeneID (NCBI):

51478

UNIPROT ID:

P56937

Full Name:

hydroxysteroid (17-beta) dehydrogenase 7

Calculated MW:

38 kDa

Observed MW:

32-38 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

Applications

Tested Applications:

WB, IP, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human, mouse

Cited Species:

human, mouse

Positive Controls:

WB : HEK-293 cells, MCF-7 cells, Jurkat cells

IP : MCF-7 cells,

Notable Publications

Author	Pubmed ID	Journal	Application
Xiaoyu Tu	29728128	Reprod Biol Endocrinol	WB, IHC
Taha A Jan	34260939	Cell Rep	IF
Liel-Sarah Izchikis	39307342	Mol Cell Endocrinol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 14854-1-AP (HSD17B7 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



WB result of HSD17B7 antibody (14854-1-AP; 1:400; incubated at room temperature for 1.5 hours) with sh-Control and sh-HSD17B7 transfected HEK-293 cells.



IP result of anti-HSD17B7 (IP:14854-1-AP, 4ug; Detection:14854-1-AP 1:1000) with MCF-7 cells lysate 3200ug.