For Research Use Only

NARS Polyclonal antibody

Catalog Number: 14882-1-AP

1 Publications



Basic Information

Applications

Catalog Number: 14882-1-AP

GenBank Accession Number:

Purification Method: Antigen affinity purification

Size:

BC001687 GeneID (NCBI):

Recommended Dilutions:

150ul, Concentration: 350 µg/ml by

4677

WB 1:500-1:2000

Nanodrop and 300 µg/ml by Bradford Full Name:

asparaginyl-tRNA synthetase

IP 0.5-4.0 ug for IP and 1:500-1:1000 for WB

method using BSA as the standard;

Calculated MW: 63 kDa

63 kDa

Isotype: IgG

Observed MW:

IHC 1:50-1:500

Immunogen Catalog Number:

AG6656

Rabbit

Tested Applications:

Positive Controls: WB: K-562 cells, HepG2 cells

IHC. IP. WB. FLISA

IP: HepG2 cells,

Cited Applications: WB

Species Specificity:

human

Cited Species:

human

IHC: human liver cancer tissue,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Asparagine-tRNA ligase (NARS) belongs to the class-II aminoacyl-tRNA synthetase family. Catalyzes the attachment of asparagine to tRNA(Asn) in a two-step reaction: asparagine is first activated by ATP to form Asn-AMP and then transferred to the acceptor end of tRNA(Asn) (PubMed: 9421509). In addition to its essential role in protein synthesis, acts as a signaling molecule that induced migration of CCR3-expressing cells. The NARS levels were higher in tumor tissues compared with adjacent normal tissues, and positively associated with LN metastasis. NARS may serve as a potential biomarker for lung adenocarcinoma (ADC) diagnosis/prognosis (PMID: 27161446).

Notable Publications

Author **Pubmed ID** Journal Application Mirjana Malnar Črnigoj Nat Commun WB 37717009

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



K-562 cells were subjected to SDS PAGE followed by western blot with 14882-1-AP (NARS antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 14882-1-AP (NARS antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP Result of anti-NARS (IP:14882-1-AP, 4ug; Detection:14882-1-AP 1:500) with HepG2 cells lysate 4000ug.