## For Research Use Only

# QDPR Polyclonal antibody

Catalog Number: 14908-1-AP

3 Publications



**Basic Information** 

Catalog Number: 14908-1-AP

GeneID (NCBI):

150ul, Concentration: 500 µg/ml by Nanodrop and 333 µg/ml by Bradford Full Name:

method using BSA as the standard;

Rabbit Isotype: IgG

Immunogen Catalog Number:

AG6705

Size:

GenBank Accession Number:

BC000576

5860

quinoid dihydropteridine reductase

Calculated MW: 26 kDa

Observed MW: 28-30 kDa

Antigen affinity purification Recommended Dilutions:

**Purification Method:** 

WB 1:500-1:2000 IP 0.5-4.0 ug for IP and 1:500-1:2000

for WB IHC 1:20-1:200

**Applications** 

**Tested Applications:** 

IHC, IP, WB, ELISA

**Cited Applications:** 

**Species Specificity:** human, mouse, rat **Cited Species:** human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: mouse liver tissue, human liver tissue

IP: mouse liver tissue.

IHC: human ovary tumor tissue,

# **Background Information**

Dihydropteridine reductase (QDPR), also named as DHPR and HDHPR, is an essential enzyme in the hydroxylating system of the aromatic amino acids, since it catalyses the regeneration of tetrahydrobiopterin (BH4), the natural cofactor of phenylalanine, tyrosine, and tryptophan hydroxylases, from the quininoid-dihydrobiopterin produced in these coupled reactions(PMID:8326489). The QDPR protein is active as a dimer, with a subunit Mr of 26 kDa(PMID:7627180). This protein belongs to the short-chain dehydrogenases/reductases (SDR) family. Defects in QDPR are the cause of BH4-deficient hyperphenylalaninemia type C (HPABH4C)(PMID:11153907).

#### Notable Publications

Author	Pubmed ID	Journal	Application
Natasha L Pacheco	29090078	Mol Autism	WB
Zhiguo Li	34643893	Genes Genomics	WB
Kalina Wiatr	34220448	Front Mol Neurosci	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## **Selected Validation Data**




mouse liver tissue were subjected to SDS PAGE followed by western blot with 14908-1-AP (QDPR antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human ovary tumor using 14908-1-AP (QDPR antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human ovary tumor using 14908-1-AP (QDPR antibody) at dilution of 1:100 (under 40x lens).



IP Result of anti-QDPR (IP:14908-1-AP, 4ug; Detection:14908-1-AP 1:1000) with mouse liver tissue lysate 4000ug.