For Research Use Only

POFUT1 Polyclonal antibody

Catalog Number: 14929-1-AP

Featured Product

13 Publications



Basic Information

Catalog Number: GenBank Accession Number:

14929-1-AP BC000582

Size: GeneID (NCBI):
150ul , Concentration: 240 ug/ml by 23509

Nanodrop and 200 ug/ml by Bradford UNIPROT ID: method using BSA as the standard; Q9H488

Source: Full Name:
Rabbit protein O-fucosyltransferase 1

Isotype: Calculated MW:
IgG 44 kDa
Immunogen Catalog Number: Observed MW:
AG6710 44 kDa

Purification Method:
Antigen affinity purification
Recommended Dilutions:

WB 1:1000-1:6000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF/ICC 1:50-1:500

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity: human, mouse, rat

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HCT 116 cells, HUVEC cells

IP: HepG2 cells,

IHC: human liver cancer tissue, mouse liver tissue

IF/ICC: HUVEC cells, A431 cells

Background Information

POFUT1 (protein O-fucosyltransferase 1), also known as GDP-fucose protein O-fucosyltransferase, is an enzyme that catalyzes the reaction that attaches fucose through an O-glycosidic linkage to a conserved serine or threonine residue in EGF domains and plays a crucial role in Notch signaling.

Notable Publications

Author	Pubmed ID	Journal	Application
Yuheng Du	30250219	Cell Death Dis	WB,IHC,IF
Rene G Feichtinger	34468083	EMBO Mol Med	WB
Dandan Zhang	31601791	Cell Death Dis	WB,IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




HCT 116 cells were subjected to SDS PAGE followed by western blot with 14929-1-AP (POFUT1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 14929-1-AP (POFUT1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HUVEC cells using POFUT1 antibody (14929-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit $\lg G(H+L)$.



IP result of anti-POFUT1 (IP:14929-1-AP, 4ug; Detection:14929-1-AP 1:500) with HepG2 cells lysate 2400ug.



Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 14929-1-AP (POFUT1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 14929-1-AP (POFUT1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).