For Research Use Only

## HAI-1 Polyclonal antibody

Catalog Number:15036-1-AP





Basic Information	Catalog Number: 15036-1-AP	GenBank Accession Number: BC004140	Purification Method: Antigen affinity purification	
	Size: 150ul , Concentration: 850 ug/ml by Nanodrop and 347 ug/ml by Bradford method using BSA as the standard; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG6665	OVA3278 Full Name: serine peptidase inhibitor, Kuni 1 Calculated MW:	Recommended Dilutions: WB 1:200-1:1000 IHC 1:50-1:500 IF/ICC 1:200-1:800	
		58 kDa Observed MW: 58-70 kDa		
Applications	Tested Applications: WB, IHC, IF/ICC, ELISA		e Controls: nan placenta tissue,	
	Species Specificity: human, mouse		IHC : human prostate cancer tissue,	
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		NIH/3T3 cells,	
Storage	Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3.			
*** 20ul sizes contain 0.1% BSA	Aliquoting is unnecessary for -20 $^{\circ}$ C s	torage		

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data






human placenta tissue were subjected to SDS PAGE followed by western blot with 15036-1-AP (HAI-1 antibody at dilution of 1:300 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 15036-1-AP (HAI-1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 15036-1-AP (HAI-1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed NIH/3T3 cells using HAI-1 antibody (15036-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).