For Research Use Only

AP1S2 Polyclonal antibody

Catalog Number: 15043-1-AP



Purification Method:

WB 1:1000-1:4000

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number:

15043-1-AP

Size:

Isotype:

AG6911

GenBank Accession Number:

BC001117

GeneID (NCBI):

150ul, Concentration: 800 ug/ml by 8905

Nanodrop; **UNIPROT ID:** P56377 Rabbit

Full Name: adaptor-related protein complex 1,

IgG sigma 2 subunit Immunogen Catalog Number: Calculated MW:

19 kDa

Observed MW: 19-24 kDa

Applications

Tested Applications:

WB, ELISA

Species Specificity:

human, rat

Positive Controls:

WB: rat epididymis tissue,

Background Information

AP1S2 is a subunit of the adaptor complex AP-1. It belongs to the adaptor complexes small subunit family. Adaptor $protein \, (AP) \, complexes \, are \, cytosolic \, heterotetramers \, that \, mediate \, the \, sorting \, of \, membrane \, proteins \, in \, the \, complexes \, are \, cytosolic \, heterotetramers \, that \, mediate \, the \, sorting \, of \, membrane \, proteins \, in \, the \, cytosolic \, heterotetramers \, that \, mediate \, the \, sorting \, of \, membrane \, proteins \, in \, the \, cytosolic \, heterotetramers \, that \, mediate \, the \, sorting \, of \, membrane \, proteins \, in \, the \, cytosolic \, heterotetramers \, that \, mediate \, the \, sorting \, of \, membrane \, proteins \, in \, the \, cytosolic \, heterotetramers \, that \, mediate \, the \, sorting \, of \, membrane \, proteins \, in \, the \, cytosolic \, heterotetramers \, that \, mediate \, the \, sorting \, of \, membrane \, proteins \, in \, the \, cytosolic \, heterotetramers \, that \, mediate \, the \, cytosolic \, heterotetramers \, that \, mediate \, the \, cytosolic \, heterotetramers \, that \, cytosolic \, heterotetramers \, that \, cytosolic \, heterotetramers \, cytosolic \, heterotetram$ secretory and endocytic pathways. AP-1 is found at the cytoplasmic face of coated vesicles located at the Golgi complex, where it mediates both the recruitment of clathrin to the membrane and the recognition of sorting signals within the cytosolic tails of transmembrane receptors.

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



rat epididymis tissue was subjected to SDS PAGE followed by western blot with 15043-1-AP (AP1S2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.