### For Research Use Only

# TARS2 Polyclonal antibody

Catalog Number: 15067-1-AP

**Featured Product** 

**4 Publications** 



**Basic Information** 

Catalog Number: 15067-1-AP

GenBank Accession Number: BC000541

**Purification Method:** Antigen affinity purification

Size:

GeneID (NCBI):

Recommended Dilutions:

150ul, Concentration: 550 µg/ml by Nanodrop and 287 µg/ml by Bradford Full Name:

80222

WB 1:500-1:1000

method using BSA as the standard;

threonyl-tRNA synthetase 2,

IP 0.5-4.0 ug for IP and 1:500-1:1000

mitochondrial (putative)

for WB IHC 1:50-1:500 IF 1:20-1:200

Rabbit Isotype: IgG

Calculated MW: 81 kDa Observed MW:

Immunogen Catalog Number:

AG7074

81 kDa

**Applications** 

**Tested Applications:** 

IF, IHC, IP, WB, ELISA

**Cited Applications:** 

IF, IHC, IP, WB

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

**Positive Controls:** 

WB: HepG2 cells, HeLa cells, mouse liver tissue

IP: mouse liver tissue.

IHC: mouse heart tissue, mouse liver tissue

IF: HepG2 cells,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

## **Background Information**

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Hui Tian	35184682	Bioengineered	WB
Sung-Hoon Kim	33340489	Mol Cell	IF, WB, IP
Lida Suo	37956648	Int J Biol Markers	IHC

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


\*\*\* 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


### **Selected Validation Data**




Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 15067-1-AP (TARS2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 15067-1-AP (TARS2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



HepG2 cells were subjected to SDS PAGE followed by western blot with 15067-1-AP (TARS2 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



IP Result of anti-TARS2 (IP:15067-1-AP, 4ug; Detection:15067-1-AP 1:600) with mouse liver tissue lysate 4000ug.



Immunofluorescent analysis of (10% Formaldehyde) fixed HepG2 cells using 15067-1-AP (TARS2 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).