## For Research Use Only

## SEC61B Polyclonal antibody

Catalog Number: 15087-1-AP 20 Publications



**Basic Information** 

Catalog Number:

15087-1-AP BC001734
Size: GeneID (NCBI):

150ul , Concentration: 400 ug/ml by 10952 Nanodrop; UNIPROT ID:

Source: P60468
Rabbit Full Name:
Isotype: Sec61 beta subunit

IgG Calculated MW:
Immunogen Catalog Number: 10 kDa

Immunogen Catalog Number: 10 kDa AG7169 Observed

7169 Observed MW:

10-15 kDa

GenBank Accession Number:

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:2000-1:12000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF-P 1:50-1:500 IF/ICC 1:50-1:500

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, IF-P, IP, ELISA

Cited Applications:
WB, IHC, IF, IP
Species Specificity:
human, mouse, rat
Cited Species:

human, mouse, monkey, zebrafish

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HEK-293 cells, HeLa cells, HepG2 cells, L02 cells,

mouse liver tissue, rat liver tissue

IP: HepG2 cells,

IHC: human liver tissue, human cervical cancer tissue

IF-P: mouse pancreas tissue,
IF/ICC: HepG2 cells,

**Background Information** 

The heteromeric SEC61 complex is composed of alpha, beta and gamma subunits. Oligomers of the SEC61 complex form a transmembrane channel where proteins are translocated across and integrated into the ER membrane (PMID: 10212142). SEC61B is the beta subunit of the SEC61 complex. It has been reported that SEC61B facilitates cotranslational protein transport and interacts with the signal peptidase during translocation (PMID: 9585408).

**Notable Publications** 

Author	Pubmed ID	Journal	Application
Hideaki Morishita	31526472	Elife	IF
Suman Rimal	34663454	Acta Neuropathol Commun	WB,IP
Stefan Norlin	26438609	Diabetes	WB, IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


\*\*\* 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 15087-1-AP (SEC61B antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 15087-1-AP (SEC61B antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human liver tissue slide using 15087-1-AP (SEC61B antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




IP result of anti-SEC61B (IP:15087-1-AP, 4ug; Detection:15087-1-AP 1:500) with HepG2 cells lysate 1800ug.



Immunofluorescent analysis of (4% PFA) fixed mouse pancreas tissue using SEC61B antibody (15087-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse pancreas tissue using SEC61B antibody (15087-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated Affini Pure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using SEC61B antibody (15087-1-AP) at dilution of 1:200 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).