For Research Use Only

NNMT Polyclonal antibody

Catalog Number: 15123-1-AP

Featured Product

21 Publications



Basic Information

Catalog Number: GenBank Accession Number:

15123-1-AP BC000234
Size: GenelD (NCBI):

Nanodrop; UNIPROT ID:
Source: P40261
Rabbit Full Name:

Isotype: nicotinamide N-methyltransferase

IgG Calculated MW:
Immunogen Catalog Number: 30 kDa

AG7197 Observed MW:

30 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:1000-1:8000

IP 0.5-4.0 ug for 1.0-3.0 mg of total $\,$

protein lysate IHC 1:400-1:1600 IF/ICC 1:20-1:200

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), IP, ELISA

150ul , Concentration: 500 ug/ml by

Cited Applications: WB, IHC, IF

Species Specificity:

human, mouse, rat Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: LO2 cells, HepG2 cells, mouse liver tissue, rat liver

tissue

IP: HepG2 cells,

IHC: human stomach cancer tissue,

IF/ICC: HepG2 cells,

Notable Publications

Author	Pubmed ID	Journal	Application
Changmei Yang	36312432	ACS Omega	WB
Changqing Wu	36382559	Adv Sci (Weinh)	WB,IHC
Wanfeng Xu	35603729	Bioengineered	WB

Storage

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com


in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




IP result of anti-NNMT (IP:15123-1-AP, 4ug; Detection:15123-1-AP 1:300) with HepG2 cells lysate 2400ug.




Immunofluorescent analysis of (10% Formaldehyde) fixed HepG2 cells using 15123-1-AP (NNMT antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug Anti-Human NNMT (15123-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



LO2 cells were subjected to SDS PAGE followed by western blot with 15123-1-AP (NNMT antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 15123-1-AP (NNMT antibody) at dilution of 1:800 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).