### For Research Use Only

# GSTM3 Polyclonal antibody

Catalog Number: 15214-1-AP

**Featured Product** 

**17 Publications** 



**Basic Information** 

Catalog Number: 15214-1-AP

GenBank Accession Number: BC000088

**Purification Method:** Antigen affinity purification

Size:

GeneID (NCBI): 2947

27-29 kDa

Recommended Dilutions: WB 1:2000-1:12000

150ul, Concentration: 350 µg/ml by Nanodrop and 220 µg/ml by Bradford Full Name:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

method using BSA as the standard;

glutathione S-transferase mu 3 (brain)  $protein\ lysate$ 

IHC 1:20-1:200

Calculated MW: Rabbit 27 kDa Isotype: Observed MW:

IgG

AG7375

Immunogen Catalog Number:

**Applications** 

**Tested Applications:** 

IHC, IP, WB, ELISA

**Cited Applications:** IHC, WB

Species Specificity:

human, mouse, rat

**Cited Species:** 

human, rat, sheep, mouse, bovine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

WB: mouse testis tissue, human testis tissue, rat testis

IP: mouse testis tissue,

IHC: human testis tissue,

## **Background Information**

GSTM3 is a cytosolic enzyme involved in prostaglandin and leukotriene synthesis and in the metabolization of toxic compounds, such as chemotherapeutic drugs, insecticides, herbicides, carcinogens, and by-products of oxidative stress. GSTM3 is expressed in testis, brain, lung, and lymphocytes. GSTM3 have been reported to be dysregulated in cancers like prostate cancer and colon cancer.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Ming Xu	31775802	Part Fibre Toxicol	WB
Hanshu Wang	34647995	Endocrinology	WB
Dong Huang	34015269	Cell Metab	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 15214-1-AP (GSTM3 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human testis using 15214-1-AP (GSTM3 antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human testis using 15214-1-AP (GSTM3 antibody) at dilution of 1:100 (under 40x lens).



IP Result of anti-GSTM3 (IP:15214-1-AP, 4ug; Detection:15214-1-AP 1:300) with mouse testis tissue lysate 4000ug.