## For Research Use Only

# GSTM3 Polyclonal antibody

Catalog Number: 15214-1-AP

Featured Product

20 Publications

**UNIPROT ID:** 



**Basic Information** 

Catalog Number:

GenBank Accession Number: 15214-1-AP BC000088

GeneID (NCBI): Size:

150ul , Concentration: 220 ug/ml by

Bradford method using BSA as the

standard; P21266 Source: Full Name:

Rabbit glutathione S-transferase mu 3 (brain)

Isotype: Calculated MW: IgG 27 kDa Immunogen Catalog Number: Observed MW:

27-29 kDa

**Applications** 

**Tested Applications:** 

Cited Applications:

Species Specificity:

human, mouse, rat

**Cited Species:** 

human, mouse, rat, bovine, sheep

TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

**Purification Method:** 

Antigen affinity purification Recommended Dilutions:

WB 1:2000-1:12000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

IF/ICC 1:200-1:800

WB, IHC, IF/ICC, IP, ELISA

WB, IHC

Note-IHC: suggested antigen retrieval with

buffer pH 6.0

Positive Controls:

WB: mouse testis tissue, human testis tissue, rat testis

IP: mouse testis tissue,

IHC: mouse testis tissue, human testis tissue

IF/ICC: HepG2 cells,

# **Background Information**

GSTM3 is a cytosolic enzyme involved in prostaglandin and leukotriene synthesis and in the metabolization of toxic compounds, such as chemotherapeutic drugs, insecticides, herbicides, carcinogens, and by-products of oxidative stress. GSTM3 is expressed in testis, brain, lung, and lymphocytes. GSTM3 have been reported to be dysregulated in cancers like prostate cancer and colon cancer.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Ming Xu	31775802	Part Fibre Toxicol	WB
Hanshu Wang	34647995	Endocrinology	WB
Dong Huang	34015269	Cell Metab	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


### **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 15214-1-AP (GSTM3 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using GSTM3 antibody (15214-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 15214-1-AP (GSTM3 antibody) at dilution of 1:100 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 15214-1-AP (GSTM3 antibody) at dilution of 1:100 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-GSTM3 (IP:15214-1-AP, 4ug; Detection:15214-1-AP 1:300) with mouse testis tissue lysate 4000ug.