

For Research Use Only

TXNL1 Polyclonal antibody

Catalog Number: 15289-1-AP **3 Publications**



Basic Information

Catalog Number: 15289-1-AP	GenBank Accession Number: BC001156	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 550 ug/ml by Nanodrop and 420 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 9352	Recommended Dilutions: WB 1:500-1:2000
Source: Rabbit	UNIPROT ID: O43396	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate
Isotype: IgG	Full Name: thioredoxin-like 1	IHC 1:100-1:400
Immunogen Catalog Number: AG7481	Calculated MW: 32 kDa	IF/ICC 1:50-1:500
	Observed MW: 32 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA	Positive Controls:
Cited Applications: WB, IHC	WB : K-562 cells, HL-60 cells
Species Specificity: human, mouse, rat	IP : HL-60 cells,
Cited Species: human	IHC : human heart tissue, human thyroid cancer tissue
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	IF/ICC : RAW 264.7 cells,

Background Information

TXNL1 (Thioredoxin-like protein 1) is also named as TRP32, TXL, TXNL and belongs to the thioredoxin family. It might be involved in the cellular response to sugar starvation stress. The predicted 289-amino acid, 32.3-kD protein has 2 distinct domains: an N-terminal domain, which is 43% identical to human TXN, and a C-terminal domain, which showed no homology to other proteins in the sequence databases (PMID:9473519). TXNL1 functions as an effector of oxidants or redox sensor that couples oxidative stress to endocytosis, by converting redox changes into a specific GDI:Rab5-mediated endocytic response (PMID:17987124).

Notable Publications

Author	Pubmed ID	Journal	Application
Caiying Wang	30290847	Vet Res	WB
Pan Ni	25348020	Curr Cancer Drug Targets	WB
W Xu	24525731	Cell Death Dis	WB, IHC

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




K-562 cells were subjected to SDS PAGE followed by western blot with 15289-1-AP (TXNL1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




IP result of anti-TXNL1 (IP:15289-1-AP, 4ug; Detection:15289-1-AP 1:300) with HL-60 cells lysate 960ug.



Immunohistochemical analysis of paraffin-embedded human heart tissue slide using 15289-1-AP (TXNL1 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human heart tissue slide using 15289-1-AP (TXNL1 Antibody) at dilution of 1:200 (under 40x lens).



Immunofluorescent analysis of (4% PFA) fixed RAW 264.7 cells using 15289-1-AP (TXNL1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L).