

For Research Use Only

# STT3B Polyclonal antibody

Catalog Number: 15323-1-AP

Featured Product

15 Publications



## Basic Information

<b>Catalog Number:</b> 15323-1-AP	<b>GenBank Accession Number:</b> BC015880	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul, Concentration: 650 µg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 201595	<b>Recommended Dilutions:</b> WB 1:500-1:3000 IHC 1:50-1:500 IF 1:50-1:500
<b>Source:</b> Rabbit	<b>Full Name:</b> STT3, subunit of the oligosaccharyltransferase complex, homolog B ( <i>S. cerevisiae</i> )	
<b>IsoType:</b> IgG	<b>Calculated MW:</b> 97 kDa	
<b>Immunogen Catalog Number:</b> AG2039	<b>Observed MW:</b> 94 kDa	

## Applications

### Tested Applications:

IF, IHC, WB, ELISA

### Cited Applications:

CoIP, IP, WB

### Species Specificity:

human, mouse, rat

### Cited Species:

human, mouse

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

WB : rat brain tissue, human brain tissue, mouse liver tissue, rat liver tissue

IHC : human liver tissue, human brain tissue

IF : HepG2 cells,

## Background Information

STT3B is a component of the N-oligosaccharyl transferase (OST) enzyme which catalyzes the transfer of a high mannose oligosaccharide from a lipid-linked oligosaccharide donor to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent polypeptide chains. N-glycosylation occurs cotranslationally and the complex associates with the Sec61 complex at the channel-forming translocon complex that mediates protein translocation across the endoplasmic reticulum (ER). STT3B seems to be involved in complex substrate specificity

## Notable Publications

Author	Pubmed ID	Journal	Application
Cecilia Lopez-Sambrooks	27694802	Nat Chem Biol	WB
Hsiang-Chi Huang	34839261	EBioMedicine	WB
Chuan Hua He	32402213	Am J Respir Cell Mol Biol	WB

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 15323-1-AP (STT3B antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 15323-1-AP (STT3B antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



human brain tissue were subjected to SDS PAGE followed by western blot with 15323-1-AP (STT3B antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 15323-1-AP (STT3B Antibody) at dilution of 1:200 (under 40x lens).