

For Research Use Only

MED18 Polyclonal antibody

Catalog Number: 15352-1-AP **1 Publications**



Basic Information

Catalog Number: 15352-1-AP	GenBank Accession Number: BC002694	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 550 µg/ml by Nanodrop and 267 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 54797	Recommended Dilutions: WB 1:1000-1:6000
Source: Rabbit	Full Name: mediator complex subunit 18	
Isotype: IgG	Calculated MW: 24 kDa	
Immunogen Catalog Number: AG7598	Observed MW: 24 kDa	

Applications

Tested Applications: WB,ELISA	Positive Controls: WB : HeLa cells,
Cited Applications: WB	
Species Specificity: human, mouse, rat	
Cited Species: mouse	

Background Information

MED18, also named as Mediator of RNA polymerase II transcription subunit 18, is a 208 amino acid protein, which belongs to the Mediator complex subunit 18 family. MED18 is a component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. MED18 as a mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. MED18 is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors.

Notable Publications

Author	Pubmed ID	Journal	Application
Wen-Shuai Tang	34233190	Cell Rep	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 15352-1-AP (MED18 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.