

For Research Use Only

NOL12 Polyclonal antibody

Catalog Number: 15456-1-AP **2 Publications**



Basic Information

Catalog Number:

15456-1-AP

Size:

150ul , Concentration: 350 µg/ml by Nanodrop and 240 µg/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG7723

GenBank Accession Number:

BC002808

GeneID (NCBI):

79159

UNIPROT ID:

Q9UGY1

Full Name:

nucleolar protein 12

Calculated MW:

25 kDa

Observed MW:

32 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:6000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IF 1:20-1:200

Applications

Tested Applications:

WB, IP, IF, ELISA

Cited Applications:

WB, IF, IHC, CoIP

Species Specificity:

human, mouse, rat

Cited Species:

human, rat

Positive Controls:

WB : HeLa cells, SH-SY5Y cells, HepG2 cells, Raji cells

IP : HeLa cells,

IF : HepG2 cells,

Notable Publications

Author	Pubmed ID	Journal	Application
Stefanie Jäger	22190034	Nature	WB
Jingtao Liu	38467618	Cell Death Discov	IHC,WB,IF,CoIP

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 15456-1-AP (NOL12 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



HeLa cells were subjected to SDS PAGE followed by western blot with 15456-1-AP (NOL12 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP result of anti-NOL12 (IP:15456-1-AP, 4ug; Detection:15456-1-AP 1:300) with HeLa cells lysate 4000ug.



Immunofluorescent analysis of (10% Formaldehyde) fixed HepG2 cells using 15456-1-AP (NOL12 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).