

For Research Use Only

HOOK3 Polyclonal antibody

Catalog Number: 15457-1-AP

Featured Product

15 Publications



Basic Information

Catalog Number:

15457-1-AP

Size:

150ul, Concentration: 600 ug/ml by Nanodrop and 400 ug/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG7724

GenBank Accession Number:

BC056146

GeneID (NCBI):

84376

UNIPROT ID:

Q86VS8

Full Name:

hook homolog 3 (Drosophila)

Calculated MW:

83 kDa

Observed MW:

83 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:4000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF/ICC 1:10-1:100

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IHC, IF, IP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : rat liver tissue, HEK-293 cells, mouse kidney tissue, human liver tissue, mouse heart tissue, mouse liver tissue

IP : mouse kidney tissue,

IHC : human kidney tissue,

IF/ICC : HepG2 cells,

Notable Publications

Author	Pubmed ID	Journal	Application
Wenhuan Guo	34598900	Cytotherapy	WB
Mara A Olenick	30373907	J Cell Biol	WB
Tianyu Wu	36395215	Science	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




rat liver tissue were subjected to SDS PAGE followed by western blot with 15457-1-AP (HOOK3 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 15457-1-AP (HOOK3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 15457-1-AP (HOOK3 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of HepG2 cells, using HOOK3 antibody 15457-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



IP result of anti-HOOK3 (IP:15457-1-AP, 4ug; Detection:15457-1-AP 1:500) with mouse kidney tissue lysate 4000ug.