

For Research Use Only

NRD1 Polyclonal antibody

Catalog Number: 15630-1-AP

3 Publications



Basic Information

Catalog Number:

15630-1-AP

Size:

150ul, Concentration: 200 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG7923

GenBank Accession Number:

BC008775

GeneID (NCBI):

4898

UNIPROT ID:

O43847

Full Name:

nardilysin (N-arginine dibasic convertase)

Calculated MW:

132 kDa

Observed MW:

135-140 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IHC 1:50-1:500

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB, IF, IHC

Species Specificity:

human, mouse

Cited Species:

human, mouse

Positive Controls:

WB : K-562 cells,

IHC : mouse testis tissue, human testis tissue

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

NRD1(Nardilysin) is also named as NRD-C(N-arginine dibasic convertase). It is a zinc metalloendopeptidase of the M16 pitrilysin family that cleaves peptides at dibasic residues. The enzyme has also been found in neural tissues of mouse embryos indicating a role in neural development. Although nardilysin is primarily found in the cytosol, it has been reported that nardilysin is located on the cell surface acting as a receptor for heparinbinding EGF-like growth factor(PMID:15809022). It has 2 isoforms produced by alternative splicing.

Notable Publications

Author	Pubmed ID	Journal	Application
Yongfeng Xu	35683488	J Clin Med	IHC
Ilian A Radichev	24848066	Diabetes	IF
Ilian A Radichev	26773144	J Immunol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



K-562 cells lysates were subjected to SDS PAGE followed by western blot with 15630-1-AP (NRD1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 15630-1-AP (NRD1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).