## For Research Use Only

## OPA3 Polyclonal antibody

Catalog Number: 15638-1-AP

**Featured Product** 

6 Publications

BC005059

GenBank Accession Number:

with chorea and spastic paraplegia)



**Basic Information** 

Catalog Number: 15638-1-AP

Size: GeneID (NCBI): 150ul, Concentration: 400 µg/ml by 80207

Nanodrop and 267 µg/ml by Bradford Full Name:

method using BSA as the standard;

Rabbit Calculated MW: 179 aa, 20 kDa Isotype: IgG Observed MW: 20 kDa

Immunogen Catalog Number: AG8103

**Tested Applications:** 

IF, IHC, IP, WB, ELISA **Cited Applications:** IHC, IP, WB

Species Specificity: human, mouse, rat **Cited Species:** 

human, rat, mouse, bovine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

**Purification Method:** 

Antigen affinity purification

Recommended Dilutions: WB 1:500-1:2000

IP 0.5-4.0 ug for IP and 1:500-1:2000

optic atrophy 3 (autosomal recessive,  $\ensuremath{\text{for WB}}$ IHC 1:50-1:500 IF 1:50-1:500

**Applications** 

Positive Controls:

WB: A549 cells, human kidney tissue, mouse thymus tissue, Jurkat cells, HeLa cells, HEK-293T cells

IP: HeLa cells.

IHC: human liver tissue, mouse colon tissue

IF: HeLa cells.

**Background Information** 

The OPA3 cDNA encodes a deduced 179-amino acid protein. Northern blot analysis demonstrated a primary transcript of approximately 5.0 kb that was ubiquitously expressed, most prominently in skeletal muscle and kidney. Mutations in this gene have been shown to result in 3-methylglutaconic aciduria type III and autosomal dominant optic atrophy and cataract.

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Miao Liu	35176941	Bioengineered	WB
Ning Meng	31881642	Cancers (Basel)	IHC
Alice E Navein	27106103	Hum Mol Genet	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 15638-1-AP (OPA3 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human liver tissue slide using 15638-1-AP (OPA3 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 15638-1-AP (OPA3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-OPA3 (IP:15638-1-AP, 4ug; Detection:15638-1-AP 1:1000) with HeLa cells lysate 1240 ug.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using OPA3 antibody (15638-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).