For Research Use Only

Collagen Type XXVII Polyclonal antibody

Catalog Number: 15673-1-AP



Basic Information

Catalog Number: GenBank Accession Number:

15673-1-AP BC007696 GeneID (NCBI): Size:

150ul, Concentration: 550 ug/ml by 85301 Nanodrop and 273 ug/ml by Bradford $\,$ UNIPROT ID: method using BSA as the standard; Q8IZC6 Source: Full Name:

Rabbit collagen, type XXVII, alpha 1

Isotype Calculated MW: 1860 aa, 87 kDa Immunogen Catalog Number: Observed MW: AG8140 68-73 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500

Applications

Tested Applications: WB, IHC, ELISA

Species Specificity: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HeLa cells, HepG2 cells, mouse brain tissue,

mouse heart tissue

IHC: human placenta tissue,

Background Information

Collagen Type XXVII, also named COL27A1, it is a member of the collagen fibrillar gene family. It is located in the extrcellular matrix. The calculated molecular weight of COL27A1 is 186 kDa. This antibody can recognize the 73 kDa isoform of target. The high level of Col27a1 expression in mouse cartilage suggested the possibility that Col27a1 may be regulated in cartilage by SOX9, as are Col2a1, Col11a2, Col9a1, and Col9a2 (PMID: 15922909). Expression analyses of the mouse col27a1 gene demonstrated high expression in cartilage, the eye and ear, but also in lung and colon. It is likely that the major protein product of COL27A1, proa1(XXVII), is a component of the extracellular matrices of cartilage and these other tissues. Study of this collagen should yield insights into normal chondrogenesis, and provide clues to the pathogenesis of some chondrody splasias and disorders of other tissues in the pathogenesis of the pathowhich this gene is expressed (PMID: 12714037).

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 15673-1-AP (Collagen Type XXVII antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 15673-1-AP (Collagen Type XXVII antibody) at dilution of 1:100 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).