## For Research Use Only

## TK1 Polyclonal antibody

Catalog Number: 15691-1-AP

**Featured Product** 

9 Publications

BC007986



**Purification Method:** 

WB 1:5000-1:50000

protein lysate

IHC 1:50-1:500

IF 1:50-1:500

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

**Basic Information** 

Catalog Number: GenBank Accession Number:

GeneID (NCBI):

150ul, Concentration: 260 µg/ml by 7083

Nanodrop;

Source: thymidine kinase 1, soluble

Rabbit Calculated MW: Isotype: 25 kDa IgG Observed MW: Immunogen Catalog Number: 25-30 kDa

15691-1-AP

Positive Controls:

WB: HEK-293 cells, MCF-7 cells, SW 1990 cells

IP: HEK-293 cells,

IHC: human breast cancer tissue,

IF: U2OS cells,

**Applications** 

**Tested Applications:** IF, IHC, IP, WB, ELISA

Cited Applications: IF, IHC, WB

Species Specificity:

human, mouse, rat

**Cited Species:** human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

**Background Information** 

TK1(Thymidine kinase, cytosolic) belongs to the thymidine kinase family and is a regulatory enzyme that can interchange between its dimeric and tetrameric forms, which have low and high catalytic efficiencies, respectively. TK1 is involved in DNA synthesis and an important proliferation marker(PMID:21178264).

**Notable Publications** 

Author	Pubmed ID	Journal	Application
Morgan S Schrock	27773744	Adv Biol Regul	WB
Foyan Lian	33218277	Cancer Biol Ther	WB
Xu Jiang	31062473	J Cell Mol Med	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 15691-1-AP (TK1 antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours.




IP result of anti-TK1(IP:15691-1-AP, 4ug; Detection:15691-1-AP 1:800) with HEK-293 cells lysate 1400 ug.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 15691-1-AP (TK1 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 15691-1-AP (TK1 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed U2OS cells using TK1 antibody (15691-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).