

For Research Use Only

CCNJ Polyclonal antibody

Catalog Number: 15694-1-AP



Basic Information

Catalog Number: 15694-1-AP	GenBank Accession Number: BC043175	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 400 µg/ml by Nanodrop;	GeneID (NCBI): 54619	Recommended Dilutions: WB 1:500-1:2000
Source: Rabbit	Full Name: cyclin J	
Isotype: IgG	Calculated MW: 43 kDa	
Immunogen Catalog Number: AG8273	Observed MW: 43 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : mouse kidney tissue, rat kidney tissue
Species Specificity: human, mouse, rat	

Background Information

CCNJ (Cyclin-J) Belongs to the Cyclin family. Cyclin J is expressed in early Drosophila embryos, and it is essential for mediating intestinal epithelial cell recovery from bacterial pore-forming toxin attack in both Drosophila and mice. Upon TLR and IFN stimulation, cyclin J restrained macrophage inflammatory responses irrespective of the cell cycle control. Cyclin J interacted with CDKs and facilitated the phosphorylation of a set of CDK substrates, including FoxK1 and Drp1. Phosphorylation inhibited FoxK1-mediated expression of glycolytic genes by reducing FoxK1 nuclear localization. Cyclin J-CDK-dependent Drp1 phosphorylation also resulted in mitochondrial fission and the suppression of ROS production (PMID: 35412849).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 15694-1-AP (CCNJ antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.