For Research Use Only

## ACSL5 Polyclonal antibody Catalog Number:15708-1-AP 19 Publications




Basic Information	Catalog Number: 15708-1-AP	GenBank Accession Number: BC007985	Purification Method: Antigen affinity purification			
	Size: 150ul , Concentration: 350 ug/ml by Nanodrop; Source:	GeneID (NCBI): 51703 UNIPROT ID:	Recommended Dilutions: WB 1:500-1:2000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate			
				Rabbit	Q9ULC5 Full Name:	IHC 1:50-1:500
				lsotype: IgG	acyl-CoA synthetase long-chain family member 5	
	Immunogen Catalog Number: AG8258	Calculated MW: 76 kDa				
		Observed MW: 66-70 kDa				
	Applications	Tested Applications:	Positive Controls:			
		WB, IHC, IP, ELISA Cited Applications:	WB : mouse liver tissue, rat liver tisue IP : mouse brain tissue,			
WB, IP						
Species Specificity: human, mouse, rat		IHC : human colon cancer tissue,				
Cited Species: human, mouse, rat						
	Note-IHC: suggested antigen ( TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0	vely, antigen				
Background Information	TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0 Acyl-CoA synthetase 5, a member of lipid biosynthesis, is the only ACSL i enterocyte apoptosis.ACSL5 could pl	vely, antigen with citrate the ACSL gene family that catalyze soform that is both, located on mitoo ay a role in promoting fatty acid-inc elated disorders(PMID:20470896). A ocated on both the mitochondrial m	chondria and functionally involved in luced lipoapoptosis in hepatocytes as CSL5 is most abundant in liver, brown embrane and endoplasmic			
	TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0 Acyl-CoA synthetase 5, a member of lipid biosynthesis, is the only ACSL i enterocyte apoptosis.ACSL5 could pl important mechanism in fatty liver-r adipose tissue, and intestine and is l reticulum(PMID:20798351). It has 3 i	vely, antigen with citrate the ACSL gene family that catalyze soform that is both, located on mitoo ay a role in promoting fatty acid-inc elated disorders(PMID:20470896). A ocated on both the mitochondrial m	hondria and functionally involved in luced lipoapoptosis in hepatocytes as CSL5 is most abundant in liver, brown embrane and endoplasmic			
	TE buffer pH 9.0; (*) Alternati   retrieval may be performed w   buffer pH 6.0   Acyl-CoA synthetase 5, a member of   lipid biosynthesis, is the only ACSL i   enterocyte apoptosis.ACSL5 could pl   important mechanism in fatty liver-   adipose tissue, and intestine and is l   reticulum(PMID:20798351). It has 3 i   Author Put	vely, antigen with citrate the ACSL gene family that catalyze soform that is both, located on mitod ay a role in promoting fatty acid-inc elated disorders(PMID:20470896). A ocated on both the mitochondrial m soforms produced by alternative spl	chondria and functionally involved in luced lipoapoptosis in hepatocytes as CSL5 is most abundant in liver, brown embrane and endoplasmic icing.			
	TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0Acyl-CoA synthetase 5, a member of lipid biosynthesis, is the only ACSL i enterocyte apoptosis.ACSL5 could pL important mechanism in fatty liver-r adipose tissue, and intestine and is 1 reticulum(PMID:20798351). It has 3 iAuthorPut He LiHe Li362	the ACSL gene family that catalyzers soform that is both, located on mitor ay a role in promoting fatty acid-inc elated disorders(PMID:20470896). A located on both the mitochondrial misoforms produced by alternative spl pomed ID Journal	uced lipoapoptosis in hepatocytes as CSL5 is most abundant in liver, brown embrane and endoplasmic icing. Application			
	TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0Acyl-CoA synthetase 5, a member of lipid biosynthesis, is the only ACSL i enterocyte apoptosis.ACSL5 could pl important mechanism in fatty liver- adipose tissue, and intestine and is l reticulum(PMID:20798351). It has 3 iAuthorPut He LiHe Li362 292	the ACSL gene family that catalyze soform that is both, located on mitoo ay a role in promoting fatty acid-ind elated disorders(PMID:20470896). A ocated on both the mitochondrial m soforms produced by alternative spl pmed ID Journal 281450 iScience	chondria and functionally involved in luced lipoapoptosis in hepatocytes as CSL5 is most abundant in liver, brown embrane and endoplasmic icing. Application WB,IP			
	TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0Acyl-CoA synthetase 5, a member of lipid biosynthesis, is the only ACSL i enterocyte apoptosis.ACSL5 could pl important mechanism in fatty liver- adipose tissue, and intestine and is l reticulum(PMID:20798351). It has 3 iAuthorPut He LiHe Li362 292	the ACSL gene family that catalyze soform that is both, located on mitod ay a role in promoting fatty acid-inc elated disorders(PMID:20470896). A ocated on both the mitochondrial m soforms produced by alternative spl pmed ID Journal 281450 iScience 765047 Nat Commun	chondria and functionally involved in luced lipoapoptosis in hepatocytes as CSL5 is most abundant in liver, brown embrane and endoplasmic icing. Application WB,IP WB			
Notable Publications	TE buffer pH 9.0; (*) Alternati   retrieval may be performed w   buffer pH 6.0   Acyl-CoA synthetase 5, a member of   lipid biosynthesis, is the only ACSL i   enterocyte apoptosis.ACSL5 could pl   important mechanism in fatty liver-ra   adipose tissue, and intestine and is 1   reticulum(PMID:20798351). It has 3 i   Author Put   He Li 362   Terytty Yang Li 292   Yuan Qu 304   Storage: Storage Buffer:	the ACSL gene family that catalyze soform that is both, located on mitoo ay a role in promoting fatty acid-ind elated disorders(PMID:20470896). A ocated on both the mitochondrial m soforms produced by alternative spl pmed ID Journal 281450 iScience 765047 Nat Commun 463708 J Surg Res	chondria and functionally involved in luced lipoapoptosis in hepatocytes as CSL5 is most abundant in liver, brown embrane and endoplasmic icing. Application WB,IP WB			
Background Information Notable Publications Storage	TE buffer pH 9.0; (*) Alternati   retrieval may be performed w   buffer pH 6.0   Acyl-CoA synthetase 5, a member of   lipid biosynthesis, is the only ACSL i   enterocyte apoptosis.ACSL5 could pl   important mechanism in fatty liver-ra   adipose tissue, and intestine and is I   reticulum(PMID:20798351). It has 3 i   Author   He Li 362   Terytty Yang Li 297   Yuan Qu 304   Storage:   Store at -20°C. Stable for one year affer	the ACSL gene family that catalyzers soform that is both, located on mitoo ay a role in promoting fatty acid-inc elated disorders(PMID:20470896). A occated on both the mitochondrial m soforms produced by alternative spl omed ID Journal 281450 iScience 765047 Nat Commun 463708 J Surg Res ere shipment.	chondria and functionally involved in luced lipoapoptosis in hepatocytes as CSL5 is most abundant in liver, brown embrane and endoplasmic icing. Application WB,IP WB			


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 15708-1-AP (ACSL5 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP result of anti-ACSL5 (IP:15708-1-AP, 4ug; Detection:15708-1-AP 1:600) with mouse brain tissue lysate 4000ug.



Immunohistochemical analysis of paraffinembedded Colorectal cancer slide using 15708-1-AP (ACSL5 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded Colorectal cancer slide using 15708-1-AP (ACSL5 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).