## For Research Use Only

## SMPX Polyclonal antibody

Catalog Number: 15791-1-AP



**Basic Information** 

Catalog Number:

15791-1-AP BC005948

GeneID (NCBI): Size:

150ul , Concentration: 500 ug/ml by 23676 Nanodrop;

**UNIPROT ID:** Q9UHP9 Rabbit Full Name:

Isotype: small muscle protein, X-linked

IgG Calculated MW: Immunogen Catalog Number: 88 aa, 10 kDa

AG8497 Observed MW:

9 kDa

GenBank Accession Number:

**Applications** 

Storage

**Tested Applications:** 

WB, IHC, ELISA

Species Specificity:

human, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

**Purification Method:** Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000 IHC 1:50-1:500

Positive Controls:

WB: pig skeletal muscle tissue, human skeletal muscle tissue, human colon tissue, pig colon tissue


IHC: human kidney tissue, human brain tissue, human lung tissue, human ovary tissue, human spleen tissue, human testis tissue

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**




human skeletal muscle tissue were subjected to SDS PAGE followed by western blot with 15791-1-AP (SMPX antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 15791-1-AP (SMPX antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 15791-1-AP (SMPX antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



pig skeletal muscle tissue were subjected to SDS PAGE followed by western blot with 15791-1-AP (SMPX antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.