

For Research Use Only

HMMR-Specific Polyclonal antibody

Catalog Number: 15820-1-AP

Featured Product

15 Publications



Basic Information

Catalog Number:

15820-1-AP

Size:

150ul, Concentration: 600 ug/ml by Nanodrop and 333 ug/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_001142556

GeneID (NCBI):

3161

UNIPROT ID:

O75330

Full Name:

hyaluronan-mediated motility receptor (RHAMM)

Calculated MW:

84 kDa

Observed MW:

84 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:2000-1:10000

IHC 1:50-1:500

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), ELISA

Cited Applications:

WB, IHC, IF, IP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

Positive Controls:

WB : HepG2 cells, K-562 cells, C6 cells, T-47D cells

IHC : human tonsillitis tissue,

IF/ICC : HepG2 cells,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Hyaluronan-mediated motility receptor (HMMR), also termed CD168, was first described by Turley in murine cells. It is reported that HMMR has an extensive coiled-coil structure (CC) that contains multiple sites for interactive partners. Initially, HMMR was considered a novel hyaluronan-mediated motility receptor and a microtubule-associated spindle assembly factor. Full-length human RHAMM is an 85 kDa coiled-coil protein that occurs both in intracellular and extracellular compartments. It has highly restricted and tightly regulated expression in most normal tissues, but is one of a number of oncogenic proteins that are exported to the cell surface in response to tissue stress by unconventional transport mechanisms. (PMID: 36750558, PMID: 30249497)

Notable Publications

Author	Pubmed ID	Journal	Application
Vishwanatha K Rao	30317586	J Cell Physiol	WB
Tianyu Wu	36395215	Science	IF
Fan Zhou	27225119	Nature	FC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 15820-1-AP (HMMR-Specific antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




1X10⁶ HepG2 cells were intracellularly stained with 0.2 ug Anti-Human HMMR-Specific (15820-1-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 15820-1-AP (HMMR-Specific Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 15820-1-AP (HMMR-Specific Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using HMMR-Specific antibody (15820-1-AP) at dilution of 1:400 and Multi-rAb CoraLite® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).