# For Research Use Only

# PKM1-specific Polyclonal antibody

Catalog Number: 15821-1-AP

Featured Product

74 Publications



### **Basic Information**

Catalog Number: GenBank Accession Number:

15821-1-AP NM\_182470 Size: Genel D (NCBI):

150ul, Concentration: 453 µg/ml by 5315 Nanodrop:

Source: PKM pyruvate kinase, muscle

Rabbit Calculated MW:
Isotype: 58 kDa
IgG Observed MW:

58 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:1000-1:4000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

# **Applications**

Tested Applications:

IHC, IP, WB, ELISA
Cited Applications:
CoIP, IF, IHC, IP, WB
Species Specificity:
human, mouse, rat
Cited Species:

human, rat, mouse, monkey, bovine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### Positive Controls:

WB: mouse skeletal muscle tissue, mouse brain tissue, rat brain tissue, mouse colon tissue, multi-cells/tissue, mouse muscle/liver tissues, mouse heart tissue, 293 cell. HepG2/MCF7 cells

IP: mouse brain tissue,

IHC: human gliomas tissue, human skeletal muscle tissue, human brain tissue, human heart tissue, mouse heart tissue

# **Background Information**

PKM, also named as OIP3, PK2, PK3, PKM, p58, THBP1, CTHBP and Tumor M2-PK, belongs to the pyruvate kinase family. It is glycolytic enzyme that catalyzes the transfer of a phosphoryl group from phosphoenolpyruvate (PEP) to ADP, generating ATP. It stimulates POU5F1-mediated transcriptional activation. PKM plays a general role in caspase independent cell death of tumor cells. PKM has 2 isoforms named PKM1 and PKM2. The primary pyruvate kinase isoform before tumour development is PK-M1; however, the primary isoform from four independent tumours is PK-M2 (PMID:18337823). This antibody is specific to PKM1 isoform.

# **Notable Publications**

Author	Pubmed ID	Journal	Application
Stefano Miglietta	36287116	Noncoding RNA	WB
Во Ма	30235220	PLoS One	WB
Di Huang	30224822	Nat Immunol	

# Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffe

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

# Selected Validation Data




mouse skeletal muscle tissue were subjected to SDS PAGE followed by western blot with 15821-1-AP (PKM1-specific antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 15821-1-AP (PKM1-specific antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 15821-1-AP (PKM1-specific antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 15821-1-AP (PKM1-specific antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 15821-1-AP (PKM1-specific antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours.



IP result of anti-PKM1-specific(IP:15821-1-AP, 4ug; Detection:15821-1-AP 1:40000) with mouse brain tissue lysate 1680 ug.



Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 15821-1-AP (PKM1-specific antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).