

# TDO2 Polyclonal antibody

Catalog Number: 15880-1-AP

Featured Product

32 Publications

## Basic Information

<b>Catalog Number:</b> 15880-1-AP	<b>GenBank Accession Number:</b> BC005355	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul, Concentration: 350 µg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 6999	<b>Recommended Dilutions:</b> WB 1:500-1:1000 IHC 1:100-1:400 IF 1:200-1:800
<b>Source:</b> Rabbit	<b>Full Name:</b> tryptophan 2,3-dioxygenase	
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 406 aa, 48 kDa	
<b>Immunogen Catalog Number:</b> AG8642	<b>Observed MW:</b> 40-50 kDa	

## Applications

<b>Tested Applications:</b> IF, IHC, WB, ELISA	<b>Positive Controls:</b>
<b>Cited Applications:</b> FC, IF, IHC, WB	<b>WB:</b> mouse liver tissue, rat liver tissue
<b>Species Specificity:</b> human, mouse, rat	<b>IHC:</b> human liver cancer tissue, human liver tissue
<b>Cited Species:</b> human, chicken, rat, mouse	<b>IF:</b> A431 cells,
<b>Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b>	

## Background Information

TDO2 (tryptophan 2,3-dioxygenase), IDO1 (indoleamine 2,3-dioxygenase 1) and IDO2 are three enzymes that catalyze the amino acid tryptophan into kynurenine in the kynurenine pathway (PMID: 23090118, 28469468). TDO2 plays an important role in neurological diseases such as Alzheimer's disease, Parkinson's disease and autism. Overexpression of TDO2 promoted tumor cell survival and was correlated with tumor grade and poor prognosis in triple negative breast cancer, brain tumors and esophageal squamous cell carcinoma. TDO2 may be a potential therapeutic target in some cancers. (PMID: 21976023, 30134247)

## Notable Publications

Author	Pubmed ID	Journal	Application
Susu Li	34529208	Transgenic Res	WB,IHC,FC,IF
Victoria L Wyckelsma	34573075	Antioxidants (Basel)	WB
Mizuho Sato	36325028	Int J Tryptophan Res	WB

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
**Aliquoting is unnecessary for -20°C storage**

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 15880-1-AP (TDO2 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Methanol) fixed A431 cells using TDO2 antibody (15880-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 15880-1-AP (TDO2 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 15880-1-AP (TDO2 Antibody) at dilution of 1:200 (under 40x lens).