For Research Use Only

KIF20A Polyclonal antibody

Catalog Number:15911-1-AP

Featured Product 11 Publications





Basic Information	Catalog Number: 15911-1-AP	GenBank Accession Number: BC012999	Purification Method: Antigen affinity purification
	Size:	GenelD (NCBI):	Recommended Dilutions:
	150ul , Concentration: 350 ug/ml by	10112	WB 1:500-1:2000
	Nanodrop and 193 ug/ml by Bradford method using BSA as the standard;	UNIFROTID.	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate
	6	095235	IHC 1:50-1:500
	Source: Rabbit	Full Name:	
	Isotype:	kinesin family member 20A Calculated MW:	
	IgG	890 aa, 100 kDa	
	Immunogen Catalog Number: AG8700	Observed MW: 100 kDa	
Applications	Tested Applications:	Positive	Controls:
	WB, IP, IHC, ELISA	WB: Cad	co-2 cells, HeLa cells
	Cited Applications: WB, IHC, IF, CoIP	IP : HEK-	-293 cells,
	Species Specificity: human	IHC : hu tissue	man breast cancer tissue, human liver cance
	Cited Species: human, rat		
	Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternativ	vely, antigen	
	retrieval may be performed w buffer pH 6.0	ith citrate	
Background Information	buffer pH 6.0 KIF20A, also known as RAB6KIFL or M involved in protein transport at the G accumulate in mitotic cells and is rec	Klp2, is a kinesin originally ider olgi apparatus. KIF20A belongs t quired for normal cell division. KI 20A is almost undetectable whil	o a large family of motor proteins that IF20A is thought to be involved in e it is overexpressed in various cancers.
	buffer pH 6.0 KIF20A, also known as RAB6KIFL or M involved in protein transport at the G accumulate in mitotic cells and is rec carcinogenesis. In normal tissues KIF KIF20A has been identified as a nove	Klp2, is a kinesin originally ider olgi apparatus. KIF20A belongs t quired for normal cell division. KI 20A is almost undetectable whil	F20A is thought to be involved in e it is overexpressed in various cancers.
	buffer pH 6.0 KIF20A, also known as RAB6KIFL or M involved in protein transport at the G accumulate in mitotic cells and is rec carcinogenesis. In normal tissues KIF KIF20A has been identified as a nove	Klp2, is a kinesin originally ider olgi apparatus. KIF20A belongs t juired for normal cell division. KI 20A is almost undetectable whil l promising candidate for antica	to a large family of motor proteins that IF20A is thought to be involved in e it is overexpressed in various cancers. ncer immunotherapy.
	buffer pH 6.0KIF20A, also known as RAB6KIFL or M involved in protein transport at the G accumulate in mitotic cells and is rec carcinogenesis. In normal tissues KIF KIF20A has been identified as a noveAuthorPutTianyu Wu363	Klp2, is a kinesin originally ider olgi apparatus. KIF20A belongs t juired for normal cell division. Ki 20A is almost undetectable whil l promising candidate for antica pmed ID Journal	to a large family of motor proteins that IF20A is thought to be involved in e it is overexpressed in various cancers. ncer immunotherapy. Application IF
	buffer pH 6.0 KIF20A, also known as RAB6KIFL or M involved in protein transport at the G accumulate in mitotic cells and is rec carcinogenesis. In normal tissues KIF KIF20A has been identified as a nove Author Put Tianyu Wu 363 Suwei Zhang 317	Klp2, is a kinesin originally ider olgi apparatus. KIF20A belongs t quired for normal cell division. KI 20A is almost undetectable whil l promising candidate for antica omed ID Journal 395215 Science	to a large family of motor proteins that IF20A is thought to be involved in e it is overexpressed in various cancers. ncer immunotherapy. Application IF
	buffer pH 6.0 KIF20A, also known as RAB6KIFL or M involved in protein transport at the G accumulate in mitotic cells and is rec carcinogenesis. In normal tissues KIF KIF20A has been identified as a nove Author Put Tianyu Wu 363 Suwei Zhang 317	Klp2, is a kinesin originally ider olgi apparatus. KIF20A belongs t juired for normal cell division. Kl 20A is almost undetectable whil l promising candidate for antica omed ID Journal 395215 Science 714645 J Biochem Mol	to a large family of motor proteins that IF20A is thought to be involved in e it is overexpressed in various cancers. ncer immunotherapy. Application IF Toxicol
Notable Publications	buffer pH 6.0 KIF20A, also known as RAB6KIFL or M involved in protein transport at the G accumulate in mitotic cells and is rec carcinogenesis. In normal tissues KIF KIF20A has been identified as a nove Author Put Tianyu Wu 363 Suwei Zhang 317 Stéphanie Miserey-Lenkei 290 Storage: Storage Buffer: PBS with 0.02% sodium azide and 50	Klp2, is a kinesin originally ider olgi apparatus. KIF20A belongs t juired for normal cell division. KI 20A is almost undetectable whil l promising candidate for antica omed ID Journal 395215 Science 714645 J Biochem Mol 093437 Nat Commun er shipment.	to a large family of motor proteins that IF20A is thought to be involved in e it is overexpressed in various cancers. ncer immunotherapy. Application IF Toxicol
Background Information Notable Publications Storage	buffer pH 6.0 KIF20A, also known as RAB6KIFL or M involved in protein transport at the G accumulate in mitotic cells and is rec carcinogenesis. In normal tissues KIF KIF20A has been identified as a nove Author Put Tianyu Wu 363 Suwei Zhang 317 Stéphanie Miserey-Lenkei 290 Storage: Storage Buffer:	Klp2, is a kinesin originally ider olgi apparatus. KIF20A belongs t juired for normal cell division. KI 20A is almost undetectable whil l promising candidate for antica omed ID Journal 395215 Science 714645 J Biochem Mol 093437 Nat Commun er shipment.	to a large family of motor proteins that IF20A is thought to be involved in e it is overexpressed in various cancers. ncer immunotherapy. Application IF Toxicol

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data






Caco-2 cells were subjected to SDS PAGE followed by western blot with 15911-1-AP (KIF20A antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.


Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 15911-1-AP (KIF20A antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 15911-1-AP (KIF20A antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-KIF20A (IP:15911-1-AP, 4ug; Detection:15911-1-AP 1:500) with HEK-293 cells lysate 1200ug.



WB result of KIF20A antibody (15911-1-AP; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-KIF20A transfected HeLa cells.