For Research Use Only

KIF20A Polyclonal antibody

Catalog Number: 15911-1-AP

Featured Product

10 Publications

BC012999

GenBank Accession Number:



Basic Information

Catalog Number: 15911-1-AP

GeneID (NCBI): Size:

150ul, Concentration: 350 µg/ml by 10112 Nanodrop and 193 µg/ml by Bradford Full Name:

method using BSA as the standard; kinesin family member 20A

Calculated MW: Rabbit 890 aa, 100 kDa Isotype: Observed MW: IgG 100 kDa

Immunogen Catalog Number:

AG8700

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:500-1:2000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

Applications

Tested Applications:

IHC, IP, WB, ELISA

Cited Applications: CoIP, IF, IHC, WB

Species Specificity:

human **Cited Species:** human, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: Caco-2 cells, HeLa cells

IP: HEK-293 cells.

IHC: human breast cancer tissue, human liver cancer

Background Information

KIF20A, also known as RAB6KIFL or MKlp2, is a kinesin originally identified as a binding partner for the rab6 GTPase involved in protein transport at the Golgi apparatus. KIF20A belongs to a large family of motor proteins that accumulate in mitotic cells and is required for normal cell division. KIF20A is thought to be involved in carcinogenesis. In normal tissues KIF20A is almost undetectable while it is overexpressed in various cancers. KIF20A has been identified as a novel promising candidate for anticancer immunotherapy.

Notable Publications

Author	Pubmed ID	Journal	Application
Tianyu Wu	36395215	Science	IF
Suwei Zhang	31714645	J Biochem Mol Toxicol	
Stéphanie Miserey-Lenkei	29093437	Nat Commun	WB,IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


*** 20ul sizes contain 0.1% BSA

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Caco-2 cells were subjected to SDS PAGE followed by western blot with 15911-1-AP (KIF20A antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 15911-1-AP (KIF20A antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 15911-1-AP (KIF20A antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP Result of anti-KIF20A (IP:15911-1-AP, 4ug; Detection:15911-1-AP 1:500) with HEK-293 cells lysate 1200ug.



WB result of KIF20A antibody (15911-1-AP; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-KIF20A transfected HeLa cells.