

For Research Use Only

# AMACR/p504S Polyclonal antibody

Catalog Number: 15918-1-AP

5 Publications



## Basic Information

### Catalog Number:

15918-1-AP

### Size:

150ul, Concentration: 400 ug/ml by Nanodrop;

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG8720

### GenBank Accession Number:

BC009471

### GeneID (NCBI):

23600

### UNIPROT ID:

Q9UHK6

### Full Name:

alpha-methylacyl-CoA racemase

### Calculated MW:

382 aa, 42 kDa

### Observed MW:

44 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB 1:1000-1:5000

IHC 1:20-1:200

## Applications

### Tested Applications:

WB, IHC, ELISA

### Cited Applications:

WB, IHC, IF

### Species Specificity:

human

### Cited Species:

human

### Positive Controls:

WB : HEK-293 cells, LNCaP cells, MCF-7 cells

IHC : human prostate cancer tissue, human renal cell carcinoma tissue

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Background Information

AMACR(Alpha-methyl acyl-CoA racemase) belongs to the CaiB/BaiF CoA-transferase family. It is a mitochondrial and peroxisomal enzyme that catalyzes the conversion of 2R stereoisomers of phytanic and pristanic acid to their S counterparts. AMACR has previously been shown to be a highly sensitive marker for colorectal and clinically localized prostate cancer (PCa). However, AMACR expression is down-regulated at the transcript and protein level in hormone-refractory metastatic PCa, suggesting a hormone-dependent expression of AMACR(PMID:12213712). It has 3 isoforms produced by alternative splicing and it can form a dimer of 70-75 kDa(PMID:21812041). Defects in AMACR are the cause of alpha-methyl acyl-CoA racemase deficiency (AMACRD) and congenital bile acid synthesis defect type 4 (CBAS4).

## Notable Publications

Author	Pubmed ID	Journal	Application
Daijun Yang	32724384	Oncol Lett	IHC
Xiqi Ding	35238066	Andrologia	IHC
Zhao-Ming Xiao	34249931	Front Cell Dev Biol	IHC

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 15918-1-AP (AMACR/p504S antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human prostate cancer using 15918-1-AP (AMACR/p504S antibody) at dilution of 1:50 (under 10x lens). Heat mediated antigen retrieved with Citric acid buffer, pH6.0.



Immunohistochemical analysis of paraffin-embedded human prostate cancer using 15918-1-AP (AMACR/p504S antibody) at dilution of 1:50 (under 40x lens). Heat mediated antigen retrieved with Citric acid buffer, pH6.0.