For Research Use Only

TELO2 Polyclonal antibody

Catalog Number: 15975-1-AP

Featured Product

16 Publications



Purification Method:

WB 1:1000-1:8000

IHC 1:50-1:500

IF 1:10-1:100

for WB

Antigen affinity purification

IP 0.5-4.0 ug for IP and 1:500-1:1000

Recommended Dilutions:

Basic Information

Catalog Number: 15975-1-AP BC017188

150ul, Concentration: 500 µg/ml by 9894

Nanodrop and 333 µg/ml by Bradford Full Name:

method using BSA as the standard;

Rabbit

IgG Immunogen Catalog Number:

Isotype:

GenBank Accession Number:

GeneID (NCBI):

TEL2, telomere maintenance 2,

homolog (S. cerevisiae)

Calculated MW: 837 aa, 92 kDa

Observed MW:

92 kDa

AG8763

Positive Controls:

WB: A431 cells, Y79 cells

IP: A431 cells.

IHC: human testis tissue,

IF: HepG2 cells,

Applications

Tested Applications: IF, IHC, IP, WB, ELISA

Cited Applications:

IP. WB

Species Specificity:

human

Cited Species: human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Background Information

TELO2 gene encodes a telomere length regulation protein TEL2 homolog. TELO2 may be involved in telomere length regulation and can form TTT complex with TTI1 and TTI2. TTT complex is required to stabilize protein levels of PIKK family proteins and is involved in the cellular resistance to DNA damage stresses, like ionizing radiation (IR), ultraviolet (UV) and MMC. The activity of mTORC1 and mTORC2 complexes, which regulate cell growth and survival in response to nutrient and hormonal signals, can be promoted, stabilized and maintained by TELO2.

Notable Publications

Author	Pubmed ID	Journal	Application
Youngran Kim	34838521	J Mol Biol	IP
Miguel Baena	25463011	Biochim Biophys Acta	WB
Jing You	27132593	Am J Hum Genet	WB

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 15975-1-AP (TELO 2 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




IP result of anti-TELO2(IP:15975-1-AP, 4ug; Detection:15975-1-AP 1:500) with A431 cells lysate 1280 ug.



Immunohistochemical analysis of paraffinembedded human testis tissue slide using 15975-1-AP (TELO2 antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human testis tissue slide using 15975-1-AP (TELO2 antibody) at dilution of 1:100 (under 40x lens).



Immunofluorescent analysis of HepG2 cells using 15975-1-AP (TELO2 antibody) at dilution of 1:25 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).