For Research Use Only

SRP19 Polyclonal antibody

Catalog Number: 16033-1-AP

4 Publications



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method: Antigen affinity purification

16033-1-AP Size:

GeneID (NCBI):

Recommended Dilutions:

150ul, Concentration: 350 µg/ml by

6728

BC010947

WB 1:500-1:2000

for WB

IF 1:20-1:200

Nanodrop and 253 $\mu g/ml$ by Bradford Full Name:

signal recognition particle 19kDa

IP 0.5-4.0 ug for IP and 1:500-1:1000

method using BSA as the standard;

Calculated MW:

Isotype:

Rabbit

144 aa, 16 kDa

Observed MW:

IgG

18-25 kDa

Immunogen Catalog Number:

AG8903

Applications

Tested Applications:

IF, IP, WB, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse, rat

Cited Species:

human

Positive Controls:

WB: human liver tissue, A549 cells, HeLa cells, K-562 cells, mouse kidney tissue, mouse liver tissue, mouse

ovary tissue, Raji cells

IP: mouse kidney tissue,

IF: HeLa cells.

Background Information

The signal recognition particle (SRP) is one of the few functional small RNP particles. The SRP couples the synthesis of membrane and secretory proteins across or into the endoplasmic reticulum (ER) membrane in eukaryotes, as well as across the bacterial plasma membrane, and chloroplast thylakoid membranes. The mammalian SRP is composed as a cross the bacterial plasma membrane, and chloroplast thylakoid membranes. The mammalian SRP is composed as a cross the bacterial plasma membrane, and chloroplast thylakoid membranes. The mammalian SRP is composed as a cross the bacterial plasma membrane, and chloroplast thylakoid membranes. The mammalian SRP is composed as a cross the bacterial plasma membrane, and chloroplast thylakoid membranes. The mammalian SRP is composed as a cross the bacterial plasma membrane is a composed as a cross the bacterial plasma membrane is a composed as a cross the bacterial plasma membrane is a composed as a cross the bacterial plasma membrane is a composed as a cross that the bacterial plasma membrane is a cross that the bacterial plasma mof a 7S (or 7SL) RNA and six different proteins, SRP9, SRP14, SRP19, SRP54, SRP68 and SRP72. All of the components of SRP, including SRP RNA, participate directly in the overall protein targeting process. SRP19 binds directly to 7S RNA and mediates binding of the 54 kDa subunit of the SRP. SRP19 was shown to significantly enhance SRP54 attachment to helix 8 of 7SL RNA. Binding of SRP19 leads to restructuring of both helix 6 and 8, causing local changes at the SRP54-binding site. This antibody is a rabbit polyclonal antibody raised against full length SRP19 of human

Notable Publications

Author	Pubmed ID	Journal	Application
Joseph Russo	28129347	PLoS One	WB
Anne-Sophie Gribling-Burrer	28115638	Nucleic Acids Res	WB
Diego Acosta-Alvear	30582518	Elife	WB

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



human liver tissue were subjected to SDS PAGE followed by western blot with 16033-1-AP (SRP19 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP Result of anti-SRP19 (IP:16033-1-AP, 4ug: Detection:16033-1-AP 1:500) with mouse kidney tissue lysate 4000ug.



Immunofluorescent analysis of HeLa cells using 16033-1-AP (SRP19 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).