

For Research Use Only

# BBOX1 Polyclonal antibody

Catalog Number: 16099-1-AP

Featured Product

2 Publications



## Basic Information

### Catalog Number:

16099-1-AP

### Size:

150ul, Concentration: 500 ug/ml by Nanodrop and 360 ug/ml by Bradford method using BSA as the standard;

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG8964

### GenBank Accession Number:

BC011034

### GeneID (NCBI):

8424

### UNIPROT ID:

O75936

### Full Name:

butyrobetaine (gamma), 2-oxoglutarate dioxygenase (gamma-butyrobetaine hydroxylase) 1

### Calculated MW:

387 aa, 45 kDa

### Observed MW:

45 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB 1:500-1:1000

IF/ICC 1:10-1:100

## Applications

### Tested Applications:

WB, IF/ICC, ELISA

### Cited Applications:

WB, IHC

### Species Specificity:

human, mouse, rat

### Cited Species:

human

### Positive Controls:

WB : human brain tissue,

IF/ICC : HepG2 cells,

## Notable Publications

Author	Pubmed ID	Journal	Application
Jiazhen Huang	39181218	Biochim Biophys Acta Mol Cell Res	WB,IHC
Jing Wang	35775648	EMBO J	WB,IHC

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



human brain tissue were subjected to SDS PAGE followed by western blot with 16099-1-AP (BBOX1 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of HepG2 cells, using BBOX1 antibody 16099-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).